



Office de la propriété
intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An Agency of
Industry Canada

7CT/BE 2004/000172

*Bureau canadien
des brevets*
Certification

*Canadian Patent
Office*
Certification

La présente atteste que les documents
ci-joints, dont la liste figure ci-dessous,
sont des copies authentiques des docu-
ments déposés au Bureau des brevets.

This is to certify that the documents
attached hereto and identified below are
true copies of the documents on file in
the Patent Office.

Specification and drawings, as originally filed, with Application for Patent Serial No:
2,487,529, on November 15, 2004, by B&B BEHEER NV, assignee of Erik Buntinx, for
"Use of D4 and 5-HT_{2A} Antagonists, Inverse Agonists or Partial Agonists".

Deacy Paulucci
Agent certificateur/Certifying Officer

June 8, 2005

Date

Canada

(CIPO 68)
31-03-04

OPIC  CIPO

BEST AVAILABLE COPY

Abstract

The present invention relates to the use of compounds and compositions of compounds having D4 and 5-HT2A antagonistic, partial agonistic or inverse agonistic activity for the treatment of the underlying dysregulation of the emotional functionality of mental disorders (i.e. affect instability – hypersensitivity – hyperaesthesia – dissociative phenomena - etc).

5 The invention also relates to methods comprising administering to a patient diagnosed as having a neuropsychiatric disorder a pharmaceutical composition containing (i) compounds having D4 antagonistic, partial agonistic or inverse agonistic activity and (ii) compounds having 5-HT2A antagonistic, partial agonistic or inverse agonistic, and (iii) any known

10 medicinal compound and compositions of said compounds. The combined D4 and 5-HT2A antagonistic, partial agonistic or inverse agonistic effects may reside within the same chemical or biological compound or in two different chemical and/or biological compounds.

USE OF D4 AND 5-HT2A ANTAGONISTS, INVERSE AGONISTS OR PARTIAL AGONISTS

Field of the Invention

The invention relates to the field of neuropsychiatry. More specifically, the invention relates to the use of compounds, which have D4 and 5-HT2A antagonist, inverse agonist or partial agonist activity, for the preparation of medicaments.

Background of the invention

Conventionally, mental disorders are divided into types based on criteria sets with defining features. DSM-IV (*American Psychiatric Association*, (1993 – ISBN 0 – 89042 – 061 – 0)) is the in the art well-known golden standard of such a categorical classification. In DSM-IV, there is no assumption that each category of mental disorder is a completely discrete entity with absolute boundaries dividing it from other mental disorders or from no mental disorder. There is also no assumption that all individuals described as having the same mental disorder are alike in all important ways. Individuals sharing a diagnosis are likely to be heterogeneous even in regard to the defining features of the diagnosis. Thus, the categorical defined mental disorders as mood and anxiety disorders are having an external and even internal variable co-incidence of symptoms concerning e.g. mood, anxiety, perception, feeding, somatic sensations, sexual functions, sleep, cognitive functioning, impulse control, attention, substance use, personality, bereavement, identity, phase of life, abuse or neglect and other aspects of behavior.

In a dimensional system, clinical presentations are classified based on quantification of attributes i.e. dysfunctions rather than the assignment to categories and works best in describing phenomena that are distributed continuously and that do not have clear boundaries.

Emotion dysregulation is known as such an attribution or dysfunction that plays an important role in the development and course of mental disorders (Gross, J. J. & Munoz, R. F., 1995, *Emotion regulation and mental health*, *Clinical Psychology: Science and Practice*, 2, 151-164; Mennin, D.S., Heimberg, R. G., Turk, C. L. & Fresco, D. M., 2002, *Applying an emotion regulation framework to integrative approaches to generalized anxiety disorder*, *Clinical Psychology: Science and Practice*, 9, 85-90; Linehan, M. M., 1993, *Cognitive-behavioral treatment of borderline personality disorder*, New York, The Guilford Press; Gratz, K. L., Roemer, L., 2001 & 2004, *Multidimensional assessment of emotion regulation and dysregulation: development, factor structure, and initial validation of the Difficulties in*

Emotion Regulation Scale, Annual meeting of the Association for Advancement of Behavior Therapy, Nov. 2001 & *Journal of Psychopathology and Behavioral Assessment*, Vol. 26, No. 1, March 2004) besides behavioural and cognitive dysfunctions. D4 dopamine receptors (D4DR), almost exclusively present in the mesocortical and mesolimbic systems (O'Malley, K. L., Harmon, S., Tang, L., Todd, R. D., *The rat dopamine D4 receptor: sequence, gene structure, and demonstration of expression in the cardiovascular system*, *New Biol.*, 4, 137-46, 1992), are in the art known as modulators of emotion and cognition. D4DR agonistic activity gives a behavioural sensitisation; D4DR antagonistic activity leads to an emotion modulation (Svensson, T. H., Mathé, A. A., *Monoaminergic Transmitter Systems, Biological Psychiatry* (eds. D'Haenen, H., et al.), 45-66, 2002, John Wiley & Sons, Ltd). Data demonstrate that agonism of the dopamine D4 receptors play an important role in the induction of behavioral sensitization to amphetamine and accompanying adaptations in pre- and postsynaptic neural systems associated with the mesolimbocortical dopamine projections (D. L. Feldpausch et al.; *The Journal of Pharmacology and Experimental Therapeutics* Vol. 286, Issue 1, 497-508, July 1998).

Results suggest that the antagonisms of cortical D2 dopamine receptors are a common target of traditional and atypical antipsychotics for therapeutic action. Higher *in vivo* binding to the D2 receptors in the cortex than in the basal ganglia is suggested as an indicator of favorable profile for a putative antipsychotic compound (X. Xiberas and J.L. Martinot; *The British Journal of Psychiatry* (2001) 179: 503-508). Results show that dopamine D4 receptor antagonism in the brain does not result in the same neurochemical consequences (increased dopamine metabolism or hyperprolactinemia) observed with typical neuroleptics (Smita Patel et al., *The Journal of Pharmacology and Experimental Therapeutics* Vol. 283, Issue 2, 636-647, 1997). The selective D4 dopamine receptor antagonist L-745,870 was ineffective as an antipsychotic for the treatment of neuroleptic responsive patients with acute schizophrenia (Kramer, M.S. et al., *Arch. Gen. Psychiatry* 1997 Dec; 54(12):1080).

Finally, in the biological system, mental disorders are defined on other levels of abstraction than in the categorical and dimensional system. Structural pathology (e.g. amyloid plaques in Alzheimer Disease), etiology (e.g. HIV Dementia) and deviance from a physiological norm (e.g. reduced cerebral blood flow) are often used as indicative biological markers for a mental disorder. The underlying dysregulation of various neurotransmitter systems (glutaminergic, GABAergic, cholinergic, monoaminergic (nor-adrenergic, dopaminergic, serotonergic), etc.) is the in the art used model for the explanation of the biological

determinants of the clinical presentation of mental disturbances. It is known that the Serotonin 2A Receptor (5-HT_{2A} receptor) - which is widespread in the Central Nervous System (CNS) - has a regulating role on the dysregulation of various neuro-transmitter systems. 5-HT_{2A} agonism gives several behavioural disturbances; 5-HT_{2A} antagonism leads to a governance of mood, social behaviour, anxiety, cognitive function, stress, sleep functions, nociception, sexual functions, feeding and other aspects of behaviour (J.E. Laysen (2004) *5-HT₂ Receptors; Current Drug Targets – CNS & Neurological Disorders*, 2004, 3, 11-26).

Dysregulation of the HPA axis (hypothalamic - pituitary - adrenal axis) has frequently been reported in patients with psychiatric disorders, and is among the most robustly demonstrated neurobiological changes among psychiatric patients (D.A. Gutman and C.B. Nemeroff, *Neuroendocrinology, Biological Psychiatry* (eds. D'Haenen, H., et al), 99, 2002, John Wiley & Sons, Ltd). The resulting elevated plasma cortisol concentrations leads to an enhanced binding of serotonin for the 5-HT_{2A} receptor (E. A. Young, *Mineralocorticoid Receptor Function in Major Depression*, *Arch Gen Psychiatry*, Jan 2003; 60: 24 - 28) and thus agonism.

Additionally 5-HT_{2A} antagonism gives a des-inhibiting of the inhibitory effect of the 5-HT_{2A} receptor on (i) the 5-HT_{1A} receptor stimulation by serotonin (S. M. Stahl, *Newer Antidepressants and Mood Stabilizers, Essential Psychopharmacology*, 265 , University Press; 2 edition (June 15, 2000); ISBN: 0521646154) and on (ii) the dopamine release in the mesocortical systems (S. M. Stahl, *Classical Antidepressants, Serotonin Selective and Noradrenargic Reuptake Inhibitors, Essential Psychopharmacology*, 233 , University Press; 2 edition (June 15, 2000); ISBN: 0521646154).

Clinical or real effectiveness of psychopharma is very rare via common pooping-out; many treatment-refractory patients and up to half of patients fail to attain remission (S. M. Stahl, *Essential Psychopharmacology, Depression and Bipolar Disorders*, 151, University Press; 2 edition (June 15, 2000); ISBN: 0521646154) . Implications of not attaining remission for Mental Disorders are increased relapse rates, continuing functional impairment and increased suicide rate (S. M. Stahl, *Essential Psychopharmacology, Depression and Bipolar Disorders*, 152 , University Press; 2 edition (June 15, 2000); ISBN: 0521646154). Clinical causes of not attaining remission by the Current Psychopharmacological Compounds are inadequate early treatment, underlying emotion dysregulation (affecting instability – hypersensitivity – hyperaesthesia – dissociative phenomena, etc.) and competitive

antagonism. There is thus a growing need for a more efficient therapy and more efficient, selective and efficacious medicaments for treating mental disorders.

Summary of the invention

- 5 The present invention relates to the use of compounds and pharmaceutical compositions having D4 and 5-HT2A antagonistic, partial agonistic or inverse agonistic activity for the treatment of the underlying emotion dysregulation of mental disorders (e.g. affecting instability – hypersensitivity – hyperaesthesia – dissociative phenomena - etc.) and to methods entailing administering to a patient diagnosed as having a mental disorder a
- 10 pharmaceutical composition containing (i) compounds having specific high selective D4 and 5-HT2A antagonistic, partial agonistic or inverse agonistic activity and (ii) a known medicinal compound and/or compositions of compounds. The combined D4 and 5-HT2A antagonistic, partial agonistic or inverse agonistic effects may reside within the same chemical or biological compound.
- 15 Taken into account the above mentioned (i) rare clinical or real effectiveness of psychotropics, (ii) the governance of the features and dysfunctions responsible - in a variable co-incidentally – for the clinical state of the mental disorders by D4 dopamine receptor (D4DR) and 2A serotonin receptor (5-HT2A) antagonism and (iii) the fact that 5-HT2A antagonism gives a des-inhibiting of the inhibitory effect of the 5-HT2A receptor on (a) the 5-HT1A
- 20 receptor stimulation by serotonin and on (b) the dopamine release in the mesocortical systems, the present invention relates to the use of a compound for the preparation of a medicament for treating a disease or disorder with an underlying emotion dysregulation, characterised in that said compound has (i) a selective affinity for the Dopamine-4 (D4) receptor with a pKi value equal to or higher than 8 towards the D4 receptor and less than 8
- 25 towards other Dopamine receptors, and (ii) a selective affinity for the 5-HT2A receptor with a pKi value equal to or higher than 8 towards the 5-HT2A receptor and less than 8 towards other 5-HT receptors and wherein said compound is administered to a patient in a dose ranging between 5 and 15 mg of the active ingredient. Preferably, said compound is pipamperon.
- 30 In a preferred embodiment, in a mono therapeutic context, the invention relates to the use of a compound as defined above, preferably pipamperon, for preparing a medicament for treating a disease or disorder selected from the group comprising anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders, factitious disorders, dissociative

disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, cognitive disorders, impulse control disorders, pervasive development, attention-deficit and disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational, identity, phase of life, academic problem, problems related to abuse or neglect.

According to a further embodiment the invention relates to the use of a first compound as defined above for the preparation of a medicament for treating a mental disease or disorder with an underlying emotion dysregulation whereby a second compound is administered simultaneously with, separate from or sequential to said first compound to augment the therapeutic effect of said second compound on said disease, or to provide a faster onset of the therapeutic effect of said second compound on said disease.

The mental diseases or disorders characterized by an underlying emotion dysregulation can be grouped into subclasses as follows: (i) non-cognitive mental disorders comprising mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, attention-deficit disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problems, identity problem, phase of life problem, academic problem and problems related to abuse or neglect; (ii) cognitive diseases comprising delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders; (iii) pain disorders; and (iv) Parkinson Disease.

In a preferred embodiment, the first compound is administered daily at least one day before administering said second compound.

Preferably, said second compound is characterized by the physiological property of influencing positively the activity of the Central Nervous System.

The invention also relates to a method for preparing a compound having a selective D4 and 5-HT2A antagonist, reverse agonist or partial agonist activity comprising the following steps:

(a) measuring the selective affinity of a test compound to the D4 receptor and selecting a compound that has a pKi value equal to or greater than 8 towards the D4 receptor in respect to all the other D receptors, and measuring the selective efficacy of the selected compound to the D4 receptor and selecting a compound which is a selective antagonist, inverse agonist or partial agonist of the D4 receptor; (b) measuring the selective affinity of a test compound to the 5-HT2A receptor and selecting a compound that has a pKi value equal to or greater than 8 towards the 5-HT2A receptor in respect to all the other 5HT receptors, and measuring the selective efficacy of the selected compound to the 5-HT2A receptor and selecting a compound which is a selective antagonist, inverse agonist or partial agonist of the 5-HT2A receptor; (c) identifying a compound which is selected in (a) and (b), (d) preparing the compound identified in (c).

The invention further also relates to a compound prepared by the described method.

Detailed description of the invention

The present inventors surprisingly found that compounds which have a high selective affinity towards the 5-HT2A receptor and which, at the same time have a high selective affinity towards the dopamine-4 (D4) receptor show an improved effect in treating underlying emotion dysregulation of mental disorders.

The compounds according to the invention may be chemical or biological in nature, or may be chemically synthesised. Preferably, the compounds of the invention are provided as a pharmaceutically acceptable salt.

One example of such a compound which has both a selective affinity for the 5-HT2A receptor with a pKi value equal to or higher than 8 towards the 5-HT2A receptor and less than 8 towards other 5HT receptors, and a selective affinity for the D4 receptor with a pKi value equal to or higher than 8 towards the D4 receptor and less than 8 towards other dopamine receptors is pipamperon. Pipamperon is the conventional name given for the compound of the formula 1'-[3-(p-Fluorobenzoyl)propyl]-[1,4'-bipiperidine]-4'-carboxamide. Pipamperon is also the active ingredient of for instance the commercially available Dipiperon (Janssen, Cilag B.V).

Further, the present inventors surprisingly found that the dosage of active ingredient for pipamperon in treatment (in monotherapy as well as in combination therapy as described in more detail further) could be very low compared to conventionally used dosages. Preferred dosages which, according to the invention, have been shown to be effective for treating

D

these mental disorders, range between 5 and 15 mg per day or between 5 and 10 mg per day. More preferably, dosages of 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 mg per day are used in treatment of the diseases of the invention. In conventional pipamperon treatment, the active ingredient is available in tablets of 40 mg per tablet or in solutions of 2 mg per drop.

5 Conventional usage of high doses ranging from 40 to 360 mg is prescribed. For instance, for children up to the age of 14, doses corresponding with 2 to 6 mg per kg body weight are conventionally prescribed. The high selective affinity of pipamperon towards the 5-HT_{2A} receptor and the D₄ receptor is reflected in the low dosage which is needed for the treatment of the mental diseases listed below and also contributes to the efficacy of the treatment.

10 The mental disorders which can be treated using pipamperon in a mono therapy at such low doses are for instance anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders, factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, cognitive disorders, impulse control disorders, pervasive development, attention-deficit and disruptive behaviour disorders,
15 substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational, identity, phase of life, academic problem, problems related to abuse or neglect.

Mental disorders such as depression are commonly treated with serotonin re-uptake inhibitors. Unfortunately, however, these compounds can give rise to side effects in use.

20 Moreover, a substantial problem in most treatment of mental disorders is the non-response to selective serotonin re-uptake inhibitors (SSRIs). Also the onset of the therapeutic effect can be delayed undesirable.

A problem to be solved by the present invention is thus the provision of a more efficient therapy and efficient, highly selective and efficacious medicaments for treating mental
25 disorders.

The inventors found that, for instance, the non-response to selective serotonin re-uptake inhibitors (SSRIs) in depression may be declared by (partial) inhibition of the 5-HT_{1A} stimulation via 5-HT_{2A} stimulation. Des-inhibition thereof via 5-HT_{2A} antagonism seems to be an answer to this problem.

30 The present inventors found that a simultaneous or foregoing treatment with a compound having a high selective 5-HT_{2A} antagonist, inverse agonist or partial agonist activity, could lead to a greater response towards, for instance, SSRIs. However, not all compounds exhibiting 5-HT_{2A} antagonism are useful: competition between 5-HT_{2A} stimulation via serotonin and 5-HT_{2A} antagonism via the compound could be responsible for the lack of more efficacy of

compounds which have both a selective serotonin re-uptake inhibitory and 5-HT_{2A} antagonist profile, such as trazodone and nefazodone.

The present inventors further surprisingly found that a simultaneous or foregoing treatment with a compound having a high selective D₄ antagonist, inverse agonist or partial agonist activity in combination with a compound having a high selective 5-HT_{2A} antagonist, inverse agonist or partial agonist activity could lead to a greater response towards, for instance, SSRIs.

In this invention, the term "antagonist" refers to an interaction between chemicals in which one partially or completely inhibits the effect of the other, in particular agents having high affinity for a given receptor, but which do not activate this receptor.

In this invention, the term "inverse agonist" refers to a ligand which produces an effect opposite to that of the agonist by occupying the same receptor.

In this invention, the term "agonist" relates to an agent which both binds to a receptor and has an intrinsic effect.

In this invention, the term "partial agonist" relates to an agent with lower intrinsic activity than a full agonist, and which produces a lower maximum effect.

The present inventors found that a compound which binds to the 5-HT_{2A} receptor with a pK_i of at least 8 but for which the binding affinity, i.e. pK_i, towards other 5HT receptors is less than 8 in combination with a high selective affinity for the D₄ receptor, i.e. which bind to the D₄ receptor with a pK_i of at least 8 but for which the binding affinity, i.e. pK_i, towards other dopamine receptors is less than 8 also show such an improved effect in treatment. These effects, i.e. D₄ antagonism, inverse agonism or partial agonism and 5-HT_{2A} antagonism, inverse agonism or partial agonism, may reside in the same compound.

The term "other 5HT receptors" as used herein relate to for instance 5-HT₁ receptors (e.g. 5-HT_{1A}, 5-HT_{1B}, 5-HT_{1D}, 5-HT_{1E}, 5-HT_{1F}), 5-HT_{2B}, 5-HT_{2C}, 5-HT₆ (rat) and 5-HT₇ (rat).

By the expression "selective affinity for the 5-HT_{2A} receptor" is meant that the receptor has a higher affinity for the 5-HT_{2A} receptor than for other 5-HT receptors.

The expression "selective affinity for the D₄ receptor" means that the receptor has a higher affinity for the dopamine D₄ receptor than for other dopamine receptors.

The term "other dopamine receptors" are, for instance, D₁, D₂ and D₃ dopamine receptors.

pK_i values of test compounds for dopamine receptors as well as 5-HT_{2A} receptors can be measured using commonly known assays.

Compounds which have a selective affinity for the D4 receptor preferably have a pKi value equal to or higher than 8 towards the D4 receptor and less than 8 towards other dopamine receptors.

Preferably, the compounds of the invention which have a selective affinity for the 5-HT2A receptor (or the D4 receptor), are compounds which have a pKi value equal to or higher than 8 towards the 5-HT2A receptor and the D4 receptor, and less than 8 towards other 5-HT receptors or dopamine receptors, respectively, as can be measured, for instance by methods known in the art. For instance, the "NIMH Psychoactive Drug Screening Program (PDSP)" K_i database (<http://kiddb.cwru.edu/nimh/5http.php>), is a unique resource in the public domain which provides information on the abilities of drugs to interact with an expanding number of molecular targets. The PDSP K_i database serves as a data warehouse for published and internally-derived pKi, or affinity, values for a large number of drugs and drug candidates at an expanding number of G-protein coupled receptors, ion channels, transporters and enzymes. The PDSP internet site also provides for commonly used protocols and assays for measuring pKi values of 5-HT and dopamine receptors.

A preferred example of a compound which has both a selective affinity for the 5-HT2A receptor with a pKi value equal to or higher than 8 towards the 5-HT2A receptor and less than 8 towards other 5-HT receptors, and a selective affinity for the D4 receptor with a pKi value equal to or higher than 8 towards the D4 receptor and less than 8 towards other dopamine receptors and which is therefore useful in a combination therapy is pipamperon.

Table 1 illustrates the selective affinity of for instance pipamperon for the 5-HT2A and for the D4 receptor. In addition, Table 1 also illustrates the low or absence of affinity of pipamperon for other receptors such as the adrenergic receptors Alpha 1A, Alpha 2A, Alpha 2B, Alpha 2C, Beta 1, Beta 2, and the histamine receptor H1. As such, treating patients with pipamperon will provide for less side effects which otherwise result from simultaneous stimulation of other receptors. Therefore, and according to preferred embodiments, useful compounds according to the invention not only have a selective 5-HT2A and/or D4 affinity but also a low affinity for other receptors such as the adrenergic and histamine receptors.

The low dosage which can be used in pipamperon treatment, as already described earlier, contributes to the high selective affinity of the compound towards the 5-HT2A receptor and the D4 receptor and therefore also to the efficacy of the treatment.

The mental diseases or disorders characterized by an underlying emotion dysregulation can be grouped into subclasses as follows: (i) the non-cognitive mental disorders comprising mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual

syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, attention-deficit disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problems, identity problem, phase of life problem, academic problem and problems related to abuse or neglect; (ii) cognitive diseases comprising delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders; (iii) the pain disorders; and (iv) Parkinson Disease. In Table 5, this classification has been used for summarizing the diseases and disorders relative to known psychotropics. In Table 6, an overview of pharmacological grouping is provided, indicating the pharmacological profile numbering, the pharmacological profile, the main disease or disorder indication(s), the name of the compound, the dose range, and the company producing or selling said compound.

These diseases and their diagnosis are very clearly defined in the "Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)" published by the American Psychiatric Association. This manual sets forth diagnostic criteria, descriptions and other information to guide the classification and diagnosis of mental disorders and is commonly used in the field of neuropsychiatry. It is for instance available on the internet under: <http://www.behavenet.com/capsules/disorders/dsm4tr.htm>.

The expression "non-cognitive diseases or disorders" used in some of the embodiments of the invention comprises the following group of diseases or disorders: mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, attention-deficit disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problems, identity problem, phase of life problem, academic problem and problems related to abuse or neglect.

In other embodiments of the invention, the mental diseases or disorders that are characterized by an underlying emotion dysregulation belong to the group of pain disorders. For instance, the combination therapy with pipamperon is especially advantageous for management of acute pain in diseases such as, but not limited to, musculoskeletal diseases, rheumatoid arthritis, osteoarthritis and ankylosing spondylitis. For the classification of pain disorders, reference is also made to the DSM-IV where these disorders are clearly described in the section of somatoform disorders by way of internationally accepted diagnostic criteria.

In other embodiments of the invention, the 5-HT_{2A} receptor and/or Dopamine-4 receptor antagonist, inverse agonist or partial agonist (e.g. pipamperon) is used in treatment of patients having neuro-degenerative diseases or disorders, or related cognitive diseases or disorders. The diseases or disorders of the present invention are characterized by an underlying degeneration of the Central Nervous System (CNS), preferably selected from the group consisting of, but not limited to, neurodegenerative diseases such as Parkinson Disease, and in other embodiments of the invention, selected from the group of (related) cognitive diseases or disorders such as Alzheimer Disease.

For instance, Parkinson Disease, which is a chronic progressive nervous disease chiefly of later life, is linked to decreased dopamine production in the substantia nigra and is marked by tremor and weakness of resting muscles and by a shuffling gait. Dopamine agonists and even levodopa, widely used in Parkinson Disease, gives via a dopamine D₄ receptor stimulation psychiatric manifestations. The induced release of serotonin acts via 5-HT_{2A} stimulation as a "brake" on dopamine release (Young B.K., Camicioli R., Ganzini L., *Neuropsychiatric adverse effects of antiparkinsonian drugs. Characteristics, evaluation and treatment. Drugs Aging.* 1997 May;10(5):367-83). Because of the need of specific D₄ and 5-HT_{2A} antagonism in the treatment of Parkinson Disease with dopamine agonists and even levodopa, it seems reasonable to combine with a compound with a high selective D₄ and 5-HT_{2A} antagonism i.e. having merely no activity towards the other receptors especially the D₂ receptor because of the primary need of the relieve of the excessive burden of remaining dopaminergic neurons. Therefore, the use of the so-called atypical anti-psychotics or serotonin-dopamine antagonists (SDAs) is absolutely contra-indicated since their high affinity for the D₂ receptor. Even the use of serotonin releasing compounds such as SSRIs in the absence of an effective 5-HT_{2A} antagonism are contra-productive towards the Parkinson Disease symptoms although many Parkinson patients are in need for an antidepressant since major depression is a very common and disabling condition in this kind of patients.

The expression "(related) cognitive diseases or disorders" according to the invention comprises, the following group of diseases or disorders: delirium (F05), dementia (such as Alzheimer Disease (F00), vascular dementia (F01), dementia due to other general medical conditions (HIV disease (F02.4), head trauma (F06.8), Parkinson Disease (F02.3), Huntington Disease (F02.2), Pick Disease (F02.0), Creutzfeldt-Jacob Disease (F02.1) and other (F02.8)), substance-induced persisting dementia (F1x.6)), amnestic disorders due to a general medical condition (F06.8) or a substance-induced persisting amnestic disorder (F1x.6), mild cognitive impairment disorder (F06.7) and other cognitive disorders (F04). The above list of diseases is provided by way of example and is not intended to limit the invention.

For instance, Alzheimer Disease is a degenerative brain disease of unknown cause that is the most common form of dementia. Alzheimer Disease usually starts in late middle age or in old age as a memory loss for recent events spreading to memories for more distant events and progresses over the course of five to ten years to a profound intellectual decline characterized by dementia and personal helplessness. The disease is marked histologically by the degeneration of brain neurons especially in the cerebral cortex and by the presence of neurofibrillary tangles and plaques containing beta-amyloid. Because dopamine receptor D4 (DRD4) antagonism can inhibit the behavioral disturbances – merely aggression and confusion - caused by the degeneration of dopamine D2 receptors (*Esiri, M.M., The basis for behavioural disturbances in dementia, J. Neurol. Neurosurg. Psychiatry, 1996; 61(2):127–130.2*) accompanied with Alzheimer disease and 5-HT2A antagonism has an important boosting effect towards the effect of cholinesterase inhibitors such as used in the treatment by facilitating the affected dopamine release in the mesocortical dopamine pathways, a high selective D4/5-HT2A-antagonist would be a more preferable compound to combine with a cholinesterase inhibitor since this avoids the counteracting effect of the in the art used SDAs on the cognitive functioning by its dopamine receptor D2-antagonism.

These diseases and their diagnoses are very clearly defined in the "*International Statistical Classification of Diseases and Related Health Problems, 1989 Revision, Geneva, World Health Organization, 1992 (ICD-10)*". This manual sets forth diagnostic criteria, descriptions and other information to guide the classification and diagnosis of neurodegenerative disorders and is commonly used in the field of neurology. According to the ICD-10 classification, the cognitive disorders are classified under several classes of disorders, i.e. dispersed under categories F00 to F19 (see above: respective classification between parentheses). Following the DSM classification, however, they are grouped in one class of diseases or disorders.

The terms "treatment", "treating", and the like, as used herein include amelioration or elimination of a developed mental disease or condition once it has been established or alleviation of the characteristic symptoms of such disease or condition. As used herein these terms also encompass, depending on the condition of the patient, preventing the onset of a disease or condition or of symptoms associated with a disease or condition, including reducing the severity of a disease or condition or symptoms associated therewith prior to affliction with said disease or condition. Such prevention or reduction prior to affliction refers to administration of the compound or composition of the invention to a patient that is not at the time of administration afflicted with the disease or condition. "Preventing" also encompasses preventing the recurrence or relapse-prevention of a disease or condition or of symptoms associated therewith, for instance after a period of improvement. It should be clear that mental conditions may be responsible for physical complaints. In this respect, the term "treating" also includes prevention of a physical disease or condition or amelioration or elimination of the developed physical disease or condition once it has been established or alleviation of the characteristic symptoms of such conditions.

As used herein, the term "medicament" also encompasses the terms "drug", "therapeutic", "potion" or other terms which are used in the field of medicine to indicate a preparation with therapeutic or prophylactic effect.

The present inventors not only found that the selective 5-HT_{2A} and D₄ antagonists, inverse agonists or partial agonists have an effect in augmenting the therapeutic effect or in providing a faster onset of the therapeutic effect of a diversity of other pharmaceutical compounds, i.e. also named "second compounds" in the present invention, in the treatment of specific diseases or disorders. A few examples of other pharmaceutical compounds whose effects are augmented or where the onset of the effect is fastened upon simultaneous or fore-going treatment with a selective 5-HT_{2A} and D₄ antagonist, preferably pipamperon in a low dose, are nor-epinephrine re-uptake inhibitors, neuroleptic agents, dopamine antagonists, or compounds used for treating or alleviating musculoskeletal diseases or disorders. A further list of other pharmaceutical compounds or second compounds useful according to the invention is provided in Table 5. It should be clear, given the general applicable character of the invention, that this list of other pharmaceutical compounds is very brief and that the invention should not be restricted to the ones exemplified herein. It should be clear that in the present invention, pipamperon is never to be seen as a "second compound".

According to the invention, it thus has been found that the compounds having a selective 5-HT2A and D4 antagonist, inverse agonist or partial agonist activity as described above are useful for augmenting the therapeutic effect of a second compound on a disease.

According to another embodiment of the invention, it has also been found that the compounds having a selective 5-HT2A and D4 antagonist, inverse agonist or partial agonist activity as described above are useful for providing a faster onset of the therapeutic effect of a second compound on a disease.

From the above it should be clear that the selective 5-HT2A and D4 antagonist, inverse agonist or partial agonist is also named 'the first compound' in the embodiments of the invention.

According to the invention, when the 5-HT2A and D4 antagonist, inverse agonist or partial agonist activity reside in separate compounds, the term "composition" may be used. Compositions of the invention comprise a first element having (i) a selective affinity for the D4 receptor with a pKi value equal to or higher than 8 towards the D4 receptor and less than 8 towards other dopamine receptors, and a second element having (ii) a selective affinity for the 5-HT2A receptor with a pKi value equal to or higher than 8 towards the 5-HT2A receptor and less than 8 towards other 5-HT receptors.

The expression "the 5-HT2A and D4 antagonist, inverse agonist or partial agonist" is used herein to indicate a single compound having both activities or to indicate the composition comprising the activities in separate elements.

It should be clear that when, in the present invention, a composition of separate elements is used instead of a single compound, this composition of separate elements may be used in combination with another, i.e. a second, compound to augment the therapeutic effect of the other, i.e. the second, compound on the same or another disease.

When the 5-HT2A and D4 antagonist, inverse agonist or partial agonist or the composition comprising both elements and the second compound are administered simultaneously, the compounds or active ingredients may be present in a single pharmaceutical composition or formulation. Alternatively the compounds or active ingredients are administered in separate pharmaceutical compositions or formulations for simultaneous or separate use. The invention thus also relates to pharmaceutical compositions comprising pipamperon and a second compound of the invention and to the uses of these pharmaceutical compositions.

When the 5-HT2A and D4 antagonist, inverse agonist or partial agonist or the composition comprising both elements of the invention are administered prior to the second compound as defined, the 5-HT2A and D4 antagonist, inverse agonist or partial agonist or the composition

comprising both elements is administered at least during 1 day prior to said second compound. Preferably, the 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist (e.g. pipamperon) or the composition comprising both elements is administered for at least 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 days prior to the administration of the second compound.

5 Preferably, the 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist (e.g. pipamperon) or the composition comprising both elements is administered for at least 2, 3, 4 or 5 weeks prior to the administration of the second compound, or even for at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months prior to the administration of the second compound.

10 According to a preferred embodiment of the invention, the above described compounds or the composition comprising both elements having a 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist activity are useful for augmenting the therapeutic effect of citalopram or for providing a faster onset of the therapeutic effect of citalopram.

15 Citalopram or citalopram hydrobromide is a selective serotonin (5-hydroxytryptamine / 5-HT) re-uptake inhibitor (SSRI) and is the conventional name given for the compound of the formula (RS)-1-[3-(dimethylamino)propyl]-1-(p-fluorophenyl)-5-phthalanarbonitrile hydrobromide.

20 According to an embodiment, a daily dose of active ingredient of SSRI, preferably citalopram, ranges between 10 and 40 mg per day. Preferably, daily doses of active ingredient ranging between 20 and 30 mg per day are administered. More preferably, a daily dose of 10, 15, 20, 25, 30, 35 or 40 mg per day is administered.

25 According to another preferred embodiment of the invention, the above described compounds or the composition comprising both elements having a 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist activity are useful for augmenting the therapeutic effect of citalopram or for providing a faster onset of the therapeutic effect of fluvoxamine.

30 Fluvoxamine or fluvoxamine maleate (luvox, fevarin) is a selective serotonin (5-HT) reuptake inhibitor (SSRI) belonging to a new chemical series, the 2-aminoethyl oxime ethers of aralkylketones. It is chemically unrelated to other SSRIs and clomipramine. It is chemically designated as 5-methoxy-4'-(trifluoromethyl) valerophenone (E)-O-(2-aminoethyl) oxime maleate (1:1).

According to an embodiment, a daily dose of active ingredient of fluvoxamine maleate ranges between 100 and 300 mg per day. Preferably, daily doses of active ingredient ranging between 150 and 200 mg per day are administered. More preferably, a daily dose of 100, 150, 200, 250 or 300 mg per day is administered.

Most of the second compounds herein described are known in the art and may be used in doses according to the supplier's or physician's prescription, or may be used according to specific embodiments described herein.

Also encompassed by the invention are pro-drugs to these second compounds or active metabolites of these compounds. For instance, for risperidone it is known that, among other products, bio transformation in the liver produces 9-hydroxyrisperidone, which is of the same pharmacological activity and intensity as parent risperidone. Therefore, also 9-hydroxyrisperidone, naturally produced or chemically synthesized may be used in the methods and uses according to the invention.

The term "active metabolite" as used herein relates to a therapeutically active compound produced by the metabolism of a parent drug. Drugs administered to treat diseases are usually transformed (metabolized) within the body into a variety of related chemical forms (metabolites), some of which may have therapeutic activity (an active metabolite).

The present invention also encompasses the use of these second compounds, administered in the form of a pharmaceutically acceptable salt in admixture with a suitable pharmaceutically acceptable excipient.

To prepare the pharmaceutical compositions, comprising the compounds or the combination of the first and second compound described herein, an effective amount of the active ingredients, in acid or base addition salt form or base form, is combined in admixture with a pharmaceutically acceptable carrier, which can take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirably in unitary dosage form suitable, for administration orally, nasal, rectally, percutaneously or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs and solutions; or solid carriers such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included.

The pharmaceutical compounds for treatment are intended for parenteral, topical, oral or local administration and generally comprise a pharmaceutically acceptable carrier and an

amount of the active ingredient sufficient to reverse or prevent the bad effects of mental disorders. The carrier may be any of those conventionally used and is limited only by chemico-physical considerations, such as solubility and lack of reactivity with the compound, and by the route of administration.

- 5 Examples of pharmaceutically acceptable acid addition salts for use in the present inventive pharmaceutical composition include those derived from mineral acids, such as hydrochloric, hydrobromic, phosphoric, metaphosphoric, nitric and sulfuric acids, and organic acids, such as tartaric, acetic, citric, malic, lactic, fumaric, benzoic, glycolic, gluconic, succinic, p-toluenesulphonic acids, and arylsulphonic, for example.
- 10 The pharmaceutically acceptable excipients described herein, for example, vehicles, adjuvants, carriers or diluents, are well-known to those who are skilled in the art and are readily available to the public. It is preferred that the pharmaceutically acceptable carrier be one that is chemically inert to the active compounds and one that has no detrimental side effects or toxicity under the conditions of use.
- 15 The following formulations for oral, aerosol, parenteral, subcutaneous, intravenous, intramuscular, interperitoneal, rectal, and vaginal administration are merely exemplary and are in no way limiting. Overall, the requirements for effective pharmaceutical carriers for parenteral compositions are well known to those of ordinary skill in the art. See
- 20 Pharmaceutics and Pharmacy Practice, J.B. Lippincott Company, Philadelphia, PA, Banker and Chalmers, eds., pages 238-250, (1982), and ASHP Handbook on Injectable Drugs, Toissel, 4th ed., pages 622-630 (1986). Topical formulations, including those that are useful for transdermal drug release, are well-known to those of skill in the art and are suitable in the context of the present invention for application to skin.

- 25 Formulations suitable for oral administration require extra considerations considering the nature of the compounds and the possible breakdown thereof if such compounds are administered orally without protecting them from the digestive secretions of the gastrointestinal tract. Such a formulation can consist of (a) liquid solutions, such as an effective amount of the compound dissolved in diluents, such as water, saline, or orange juice; (b) capsules, sachets, tablets, lozenges, and troches, each containing a predetermined
- 30 amount of the active ingredient, as solids or granules; (c) powders; (d) suspensions in an appropriate liquid; and (e) suitable emulsions. Liquid formulations may include diluents, such as water and alcohols, for example, ethanol, benzyl alcohol, and the polyethylene alcohols, either with or without the addition of a pharmaceutically acceptable surfactant, suspending agent, or emulsifying agent. Capsule forms can be of the ordinary hard- or soft-shelled
- 35 gelatin type containing, for example, surfactants, lubricants, and inert fillers, such as lactose,

5 sucrose, calcium phosphate, and corn starch. Tablet forms can include one or more of
lactose, sucrose, mannitol, corn starch, potato starch, alginic acid, microcrystalline cellulose,
acacia, gelatin, guar gum, colloidal silicon dioxide, croscarmellose sodium, talc, magnesium
stearate, calcium stearate, zinc stearate, stearic acid, and other excipients, colorants,
10 diluents, buffering agents, disintegrating agents, moistening agents, preservatives, flavoring
agents, and pharmacologically compatible excipients. Lozenge forms can comprise the
active ingredient in a flavor, usually sucrose and acacia or tragacanth, as well as pastilles
comprising the active ingredient in an inert base, such as gelatin and glycerin, or sucrose
and acacia, emulsions, gels, and the like containing, in addition to the active ingredient, such
15 excipients as are known in the art.

The compounds of the present invention, alone or in combination with other suitable
components, can be made into aerosol formulations to be administered via inhalation. For
aerosol administration, the compounds are preferably supplied in finely divided form along
with a surfactant and propellant. Typical percentages of compounds are 0.01%-20% by
15 weight, preferably 1%-10%. The surfactant must, of course, be nontoxic, and preferably
soluble in the propellant. Representative of such agents are the esters or partial esters of
fatty acids containing from 6 to 22 carbon atoms, such as caproic, octanoic, lauric, palmitic,
stearic, linoleic, linolenic, olesteric and oleic acids with an aliphatic polyhydric alcohol or its
cyclic anhydride. Mixed esters, such as mixed or natural glycerides may be employed. The
20 surfactant may constitute 0.1%-20% by weight of the compounds, preferably 0.25-5%. The
balance of the compounds is ordinarily propellant. A carrier can also be included as desired,
e.g., lecithin for intranasal delivery. These aerosol formulations can be placed into
acceptable pressurized propellants, such as dichlorodifluoromethane, propane, nitrogen, and
the like. They also may be formulated as pharmaceuticals for non-pressured preparations,
25 such as in a nebulizer or an atomizer. Such spray formulations may be used to spray
mucosa.

It will be understood that, apart from daily doses, the compounds can be administered by
other schedules. For instance, the present invention also contemplates depot injection, in
which a long acting form of the active compound is injected into the body, such as the
30 muscles. From there the active compound slowly enters the rest of the body, so one injection
can last from 1 to 4 weeks or even multiple months. Other form of dosage administrations
relate to "once-a-week" pills, in which the ingredient is slowly released over a period of a
week, and slow-release patches, e.g. a CDS (Continuous Delivery System), or Once-a-Day
Transdermal Patches.

According to a further embodiment, the invention also relates to a method for preparing a compound or composition having a selective D4 and 5-HT_{2A} antagonist, reverse agonist or partial agonist. The invention also relates to the compounds prepared by the claimed method, with the proviso that said compound is not an already known compound, such as pipamperon.

It should be clear that the compounds and compositions described herein are useful for treating any patient in need thereof. As used herein the term "patient" is not restricted to humans but also to other mammals, for instance, domestic animals which may also suffer from any form of a mental disease or disorder described herein.

The second compounds of the invention can be further grouped according to their pharmacological profile, which is summarized in Table 6.

The present invention is now described in more detail by the following embodiments. The compounds belonging to different pharmacological profiles can be further grouped according to their action on the same pathway or system as follows.

1: combination therapy with a 5-HT (serotonin) reuptake enhancer

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D4 receptor, for instance pipamperon, in a combination therapy with a 5-HT (serotonin) reuptake enhancer, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt

thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT (serotonin) reuptake enhancer compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT (serotonin) reuptake enhancer compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT (serotonin) reuptake enhancer compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT (serotonin) reuptake enhancer compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said 5-HT (serotonin) reuptake enhancer compound is tianeptine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, tianeptine is to be administered in a daily dose ranging between 25 and 50 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT (serotonin) reuptake enhancer, preferably tianeptine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorder, personality disorder, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

A pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said 5-HT

(serotonin) reuptake enhancer is tianeptine, preferably provided in a unitary dose of between 25 and 50 mg of the active ingredient.

5 2: combination therapy with a 5-HT₁ autoreceptor agonist

10 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a 5-HT₁ autoreceptor agonist, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

15 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT₁ autoreceptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT₁ autoreceptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

30 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT₁ autoreceptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT₁ autoreceptor agonist compound, further characterized in that pipamperon is to be

35

administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said 5-HT₁ autoreceptor agonist compound is sunepitron or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT₁ autoreceptor agonist, preferably sunepitron or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorder, personality disorder, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

3: combination therapy with a 5-HT_{1A} (serotonin 1A receptor) agonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a 5-HT_{1A} (serotonin 1A receptor) agonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, Alzheimer Disease, substance-induced persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{1A} (serotonin 1A receptor) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{1A} (serotonin 1A receptor) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention further also relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{1A} (serotonin 1A receptor) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{1A} (serotonin 1A receptor) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention further also relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group consisting of Alzheimer Disease, substance-induced persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the

administration of a 5-HT_{1A} (serotonin 1A receptor) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{1A} (serotonin 1A receptor) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention further also relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{1A} (serotonin 1A receptor) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{1A} (serotonin 1A receptor) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said 5-HT_{1A} (serotonin 1A receptor) agonist compound is chosen from the group consisting of MN-305, zalospiroline, xaliproden, VPI-013 (also known as OPC-14523), tandospirone, sarizotan, PRX-00023, metanspiroline, lesopitron, gepirone, flesinoxan, EMD 68843, buspiroline, bupropion (preferably controlled release formulation) and alnespiroline, preferably xaliproden, sarizotan, gepirone, flesinoxan and bupropion (preferably controlled release formulation) or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said 5-HT_{1A} (serotonin 1A receptor) agonist is xaliproden and is to be administered in a daily dose ranging between 1 and 2 mg of the active ingredient. Even more preferably, said 5-HT_{1A} (serotonin 1A receptor) agonist is bupropion (controlled release formulation) and is to be administered in a daily dose ranging between 150 and 450 mg of the active ingredient. Even more preferably, said 5-HT_{1A} (serotonin 1A receptor) agonist is gepirone and is to be administered in a daily dose, ranging between 20 and 80 mg of the active ingredient per day.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT_{1A} (serotonin 1A receptor) agonist, preferably chosen from the group consisting of MN-305, zalospiroline, xaliproden, VPI-013 (also known as OPC-14523), tandospirone, sarizotan, PRX-00023, metanspiroline, lesopitron, gepirone, flesinoxan, EMD 68843, buspiroline, bupropion (preferably controlled release formulation) and alnespiroline, more

preferably xaliproden, sarizotan, gepirone, flesinoxan and bupropion (preferably controlled release formulation), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, Alzheimer Disease, substance-induced persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease.

The present invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said 5-HT_{1A} (serotonin 1A receptor) agonist is xaliproden, preferably provided in a unitary dose of between 1 and 2 mg of the active ingredient.

The present invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said 5-HT_{1A} (serotonin 1A receptor) agonist is bupropion (controlled release formulation), preferably provided in a unitary dose of between 150 and 450 mg of the active ingredient.

The present invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said 5-HT_{1A} (serotonin 1A receptor) agonist is gepirone, preferably provided in a unitary dose of between 20 and 80 mg of the active ingredient.

4: combination therapy with a 5-HT_{1A} (serotonin 1A receptor) antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a 5-

HT1A (serotonin 1A receptor) antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, adjustment disorders, impulse control disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT1A antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT1A antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said 5-HT1A antagonist compound is chosen from the group consisting of robalzotan tartrate hydrate and NAD299 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT1A antagonist, preferably chosen from the group consisting of robalzotan tartrate hydrate and NAD299, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, adjustment disorders, impulse control disorders, attention-deficit disorders,

substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect.

5: combination therapy with a 5-HT1B (serotonin 1B receptor) antagonist compound

5 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT2A and D4 receptor, for instance pipamperon, in a combination therapy with a 5-HT1B (serotonin 1B receptor) antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect.

15 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT1B antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT1B antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

30 According to a preferred embodiment, the invention relates to the use as described above, wherein said 5-HT1B antagonist compound is chosen from the group consisting of elzasonan, AZD1134 and AR-A2, preferably elzasonan, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

35 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT1B antagonist, preferably chosen from the group consisting of elzasonan, AZD1134 and AR-A2, preferably elzasonan, or a pro-drug or an active metabolite thereof, or

a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect.

6: combination therapy with a 5-HT_{2B} (serotonin 2B receptor) antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a 5-HT_{2B} (serotonin 2B receptor) antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{2B} antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{2B} antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT2B antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT2B antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said 5-HT2B antagonist compound is agomelatine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, agomelatine is to be administered in a daily dose ranging between 25 and 50 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT2B antagonist, preferably agomelatine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

A pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said 5-HT2B antagonist is agomelatine, preferably provided in a unitary dose of between 25 and 50 mg of the active ingredient.

7: combination therapy with a 5-HT2C (serotonin 2C receptor) antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT2A and D4 receptor, for instance pipamperon, in a combination therapy with a 5-HT2C (serotonin 2C receptor) antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual

syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{2C} antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{2C} antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{2C} antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{2C} antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said 5-HT_{2C} antagonist compound is chosen from the group consisting of SB 243213 and agomelatine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, agomelatine is to be administered in a daily dose ranging between 25 and 50 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT_{2C} antagonist, preferably chosen from the group consisting of SB 243213 and agomelatine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said 5-HT_{2C} antagonist is agomelatine, preferably provided in a unitary dose of between 25 and 50 mg of the active ingredient.

8: combination therapy with a 5-HT₃ (serotonin 3 receptor) antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a 5-HT₃ (serotonin 3 receptor) antagonist compound, are substance-related disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of substance-related disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT₃ antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT₃ antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said 5-HT₃ antagonist compound is ondansetron or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, ondansetron is to be administered in a daily dose ranging between 8 and 32 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT₃ antagonist, preferably ondansetron or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of substance-related disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said 5-HT₃ antagonist is ondansetron, preferably provided in a unitary dose of between 8 and 32 mg of the active ingredient.

9: combination therapy with a 5-HT₆ (serotonin 6 receptor) antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a 5-HT₆ (serotonin 6 receptor) antagonist compound, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT₆ antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT₆ antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said 5-HT6 antagonist compound is chosen from the group consisting of SB-271046, 742457 and 271046 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT6 antagonist, preferably chosen from the group consisting of SB-271046, 742457 and 271046 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders.

10: combination therapy with an acetylcholinesterase inhibitor compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an acetylcholinesterase inhibitor compound, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical

condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an acetylcholinesterase inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said acetylcholinesterase inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said acetylcholinesterase inhibitor compound is chosen from the group consisting of tacrine, rivastigmine tartrate, rivastigmine, physostigmine, phenserine tartrate, metrifonate, huperzine A, galantamine (preferably extended release formulation), donepezil, dichlorvos and anseculin hydrochloride, preferably tartrate, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, rivastigmine tartrate is to be administered in a daily dose ranging between 3 and 12 mg of the active ingredient. Preferably, phenserine tartrate is to be administered in a daily dose ranging between 20 and 30 mg of the active ingredient. Preferably, galantamine (extended release formulation) is to be administered in a daily dose ranging between 8 and 24 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an acetylcholinesterase inhibitor, preferably chosen from the group consisting of tacrine, rivastigmine tartrate, rivastigmine, physostigmine, phenserine tartrate, metrifonate, huperzine A, galantamine (preferably extended release formulation), donepezil, dichlorvos and anseculin hydrochloride, preferably tartrate, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

35

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said acetylcholinesterase inhibitor is rivastigmine tartrate, preferably provided in a unitary dose of between 3 and 12 mg of the active ingredient.

5

The invention further relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said acetylcholinesterase inhibitor is phenserine tartrate, preferably provided in a unitary dose of between 20 and 30 mg of the active ingredient.

10

In addition, the invention relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said acetylcholinesterase inhibitor is galantamine (preferably extended release formulation), preferably provided in a unitary dose of between 8 and 24 mg of the active ingredient.

15

11: combination therapy with an adenosine A2a receptor antagonist compound

The mental disorder which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an adenosine A_{2a} receptor antagonist compound, is Parkinson disease.

20

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an adenosine A_{2a} receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said adenosine A_{2a} receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

25

30

According to a preferred embodiment, the invention relates to the use as described above, wherein said adenosine A_{2a} receptor antagonist compound is KW-6002 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, KW-6002 is to be administered in a daily dose ranging between 40 and 80 mg of the active ingredient.

35

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an adenosine A2a receptor antagonist, preferably KW-6002 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson disease.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said acetylcholinesterase inhibitor is KW-6002, preferably provided in a unitary dose of between 40 and 80 mg of the active ingredient.

12: combination therapy with an adrenergic transmitter releaser

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an adrenergic transmitter releaser, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an adrenergic transmitter releaser compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said adrenergic transmitter releaser compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an adrenergic transmitter releaser compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said adrenergic transmitter releaser compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said adrenergic transmitter releaser compound is pipoxazole or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, pipoxazole is to be administered in a daily dose ranging between 30 and 60 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an adrenergic transmitter releaser, preferably pipoxazole, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said adrenergic transmitter releaser is pipoxazole, preferably provided in a unitary dose of between 30 and 60 mg of the active ingredient.

13: combination therapy with an alpha 1 adrenoreceptor antagonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an

alpha 1 adrenoreceptor antagonist, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders and Parkinson disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an alpha 1 adrenoreceptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said alpha 1 adrenoreceptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an alpha 1 adrenoreceptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said alpha 1 adrenoreceptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from

or prior to the administration of an alpha 1 adrenoreceptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said alpha 1 adrenoreceptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said alpha 1 adrenoreceptor antagonist compound is chosen from the group consisting of SDZ NVI 085 and flesinoxan or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an alpha 1 adrenoreceptor antagonist, preferably chosen from the group consisting of SDZ NVI 085 and flesinoxan or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders and Parkinson disease.

14: combination therapy with an alpha 2 adrenoreceptor antagonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an alpha 2 adrenoreceptor antagonist, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying

emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a alpha 2 adrenoreceptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said alpha 2 adrenoreceptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an alpha 2 adrenoreceptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said alpha 2 adrenoreceptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said alpha 2 adrenoreceptor antagonist compound is chosen from the group consisting of UK-14304, sunepitron, mirtazepine, idazoxan, fluparoxan, A75200 and (R)-A 75200, preferably sunepitron or idazoxan, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, idazoxan is to be administered in a daily dose ranging between 5 and 40 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an alpha 2 adrenoreceptor antagonist, preferably chosen from the group consisting of UK-14304, sunepitron, mirtazepine, idazoxan, fluparoxan, A75200 and (R)-A 75200, preferably sunepitron or idazoxan, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental

D

disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said alpha 2 adrenoreceptor antagonist is idazoxan, preferably provided in a unitary dose of between 5 and 40 mg of the active ingredient.

15: combination therapy with an AMPA receptor mediator compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an AMPA (alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionate) receptor mediator compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders,

impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an AMPA receptor mediator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said AMPA receptor mediator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an AMPA receptor mediator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said AMPA receptor mediator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an AMPA receptor mediator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said AMPA receptor mediator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said AMPA receptor mediator compound is chosen from the group consisting of

ampakine ORG 24448/CX-619, ampakine CX-717, ampakine CX-691 and ampakine CX-516, preferably ampakine ORG 24448/CX-619, ampakine CX-717 or ampakine CX-691, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

5 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and
(b) an AMPA receptor mediator, preferably chosen from the group consisting of ampakine
10 ORG 24448/CX-619, ampakine CX-717, ampakine CX-691 and ampakine CX-516,
preferably ampakine ORG 24448/CX-619, ampakine CX-717 or ampakine CX-691, or a pro-
drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a
combined preparation for simultaneous, separate or sequential use for treating the
underlying emotion dysregulation of a mental disease or disorder which is chosen from the
group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic
disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain
15 disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders,
impulse control disorders, pervasive development disorders, disruptive behaviour disorders,
substance-related disorders, personality disorders, psychological factors affecting medical
conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity
problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease,
20 substance-related persisting dementia, vascular dementia, dementia due to HIV disease,
dementia due to head trauma, dementia due to Parkinson Disease, dementia due to
Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob
Disease, amnesic disorders due to a general medical condition, substance-induced
persisting amnesic disorder, mild cognitive impairment disorder and other cognitive
disorders.

25

16: combination therapy with an amphetamine compound

The mental disorders which can be treated using compounds having a high selective affinity
for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an
amphetamine compound, are attention-deficit disorders.

30

The present invention thus relates to the use of pipamperon or a pharmaceutically
acceptable salt thereof for the preparation of a medicament for treating the underlying
emotion dysregulation of attention deficit disorders, characterized in that pipamperon or said
pharmaceutically acceptable salt thereof is administered simultaneously with, separate from
35 or prior to the administration of an amphetamine compound to augment the therapeutic
effect or to provide a faster onset of the therapeutic effect of said amphetamine compound,

further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said amphetamine compound is methylphenidate (preferably administered by the transdermal system) or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an amphetamine, preferably methylphenidate (preferably administered by the transdermal system) or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of attention deficit disorders.

17: combination therapy with an amyloid aggregation-inhibitor compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an amyloid aggregation-inhibitor compound, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an amyloid aggregation-inhibitor compound to augment the

therapeutic effect or to provide a faster onset of the therapeutic effect of said amyloid aggregation-inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5

According to a preferred embodiment, the invention relates to the use as described above, wherein said amyloid aggregation-inhibitor compound is chosen from the group consisting of APAN and Alzhemed, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, Alzhemed is to be administered in a daily dose of

10

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an amyloid aggregation-inhibitor, preferably chosen from the group consisting of APAN and Alzhemed, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to

15

20

Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

25

18: combination therapy with an androgen receptor modulator compound

30

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an androgen receptor modulator compound, are sexual and gender identity disorders.

35

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying

emotion dysregulation of sexual and gender identity disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an androgen receptor modulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said androgen receptor modulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said androgen receptor modulator compound is LGD2226 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an androgen receptor modulator, preferably LGD2226 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of sexual and gender identity disorders.

19: combination therapy with an beta 3 adrenoreceptor agonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an beta 3 adrenoreceptor agonist, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is

administered simultaneously with, separate from or prior to the administration of a beta 3 adrenoreceptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said beta 3 adrenoreceptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an beta 3 adrenoreceptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said beta 3 adrenoreceptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said beta 3 adrenoreceptor agonist compound is SR 58611 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a beta 3 adrenoreceptor agonist, preferably SR 58611 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

20: combination therapy with a calcium channel modulator compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a calcium channel modulator compound, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia,

dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a calcium channel modulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said calcium channel modulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a calcium channel modulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said calcium channel modulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said calcium channel modulator compound is chosen from the group consisting of safinamide and MEM 1003, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a calcium channel modulator, preferably chosen from the group consisting of safinamide and MEM 1003, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson disease.

21: combination therapy with a cannabiod receptor 1 (CB1) antagonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a cannabiod receptor 1 (CB1) antagonist, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a

cannaboid receptor 1 antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said cannaboid receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a cannaboid receptor 1 antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said cannaboid receptor 1 antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

10

15

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a cannaboid receptor 1 antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said cannaboid receptor 1 antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

20

25

According to a preferred embodiment, the invention relates to the uses as described above, wherein said cannaboid receptor antagonist compound is SR 141716 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

30

35

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a cannaboid receptor antagonist, preferably SR 141716 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development

disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

22: combination therapy with a cathepsin K inhibitor compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a cathepsin K inhibitor compound, are pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a cathepsin K inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said cathepsin K inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said cathepsin K inhibitor compound is 462795 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a cathepsin K inhibitor, preferably 462795 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of pain disorders.

23: combination therapy with a choline uptake enhancer compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a choline uptake enhancer compound, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due

to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

- 5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a choline uptake enhancer compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said choline uptake enhancer compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.
- 20 According to a preferred embodiment, the invention relates to the use as described above, wherein said choline uptake enhancer compound is MKC-231 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, MKC-231 is to be administered in a daily dose of between 20 and 160 mg of the active ingredient.
- 25 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a choline uptake enhancer, preferably MKC-231 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said choline uptake enhancer is MKC-231, preferably provided in a unitary dose of between 20 and 160 mg of the active ingredient.

24: combination therapy with a COX-2 inhibitor compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a COX-2 inhibitor compound, are pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a COX-2 inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said COX-2 inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said COX-2 inhibitor compound is chosen from the group consisting of valdecoxib, rofecoxib, parecoxib, etoricoxib, COX 189, celecoxib and ABT-963, preferably parecoxib, etoricoxib or COX 189, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, parecoxib is to be administered in a daily dose of between 20 and 80 mg of the active ingredient. Preferably, etoricoxib is to be administered in a daily dose of between 20 and 120 mg of the active ingredient. Preferably, COX 189 is to be administered in a daily dose of between 100 and 800 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a COX-2 inhibitor, preferably chosen from the group consisting of valdecoxib, rofecoxib, parecoxib, etoricoxib, COX 189, celecoxib and ABT-963, preferably parecoxib, etoricoxib or COX 189, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said COX-2 inhibitor is parecoxib, preferably provided in a unitary dose of between 20 and 80 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said COX-2 inhibitor is etoricoxib, preferably provided in a unitary dose of between 20 and 120 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said COX-2 inhibitor is COX 189, preferably provided in a unitary dose of between 100 and 800 mg of the active ingredient.

25: combination therapy with a COX-inhibiting nitric oxide donator (CINOD) compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a COX-inhibiting nitric oxide donator (CINOD) compound, are pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a COX-inhibiting nitric oxide donator (CINOD) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said COX-inhibiting nitric oxide donator (CINOD) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said COX-inhibiting nitric oxide donator (CINOD) compound is chosen from the group consisting of AZD4717 and AZD3582 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, AZD3582 is to be administered in a daily dose ranging between 93.75 and 750 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a COX-inhibiting nitric oxide donator (CINOD), preferably chosen from the group consisting of AZD4717 and AZD3582 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said COX-inhibiting nitric oxide donator (CINOD) is AZD3582, preferably provided in a unitary dose of between 93.75 and 750 mg of the active ingredient.

26: combination therapy with a CRF1 (corticotropin-releasing factor receptor 1) antagonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a CRF1 (Corticotropin-Releasing Factor receptor 1) antagonist, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a CRF1 antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said CRF1 antagonist compound, further characterized in that

pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a CRF1 antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said CRF1 antagonist
10 compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said CRF1 antagonist compound is chosen from the group consisting of R121919,
15 NBI-34041, elzasonan, CP-448,187, CP-154-526, AAG 561 and 723620, preferably R121919, elzasonan or AAG 561, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, R121919 is to be administered in a daily dose of between 5 and 80 mg of the active ingredient.

20 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a CRF1 antagonist, preferably chosen from the group consisting of R121919, NBI-34041, elzasonan, CP-448,187, CP-154-526, AAG 561 and 723620, preferably R121919, elzasonan or AAG 561, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for
25 treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related
30 disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient
35 and wherein said CRF1 antagonist is R121919, preferably provided in a unitary dose of between 5 and 80 mg of the active ingredient.

27: combination therapy with a D1 (dopamine 1) receptor agonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a D1 (dopamine 1) receptor agonist, are chosen from the group of diseases or disorders consisting of substance related disorders and Parkinson disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of substance related disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D1 receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D1 receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D1 receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D1 receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said D1 receptor agonist compound is DAS-431 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a D1 receptor agonist, preferably DAS-431 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of substance related disorders and Parkinson disease.

28: combination therapy with D2 (dopamine 2) receptor antagonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with D₂ (dopamine 2) receptor antagonist, are chosen from the group of diseases or disorders consisting of mood disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting mood disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D₂ receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D₂ receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D₂ receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D₂ receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D2 receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D2 receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said D2 receptor antagonist compound is chosen from the group consisting of bifeprunox, amisulpride aminosultopride and amisulpride, preferably bifeprunox, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a D2 receptor antagonist, preferably chosen from the group consisting of bifeprunox, amisulpride aminosultopride and amisulpride, preferably bifeprunox, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

29: combination therapy with D3 (dopamine 3) receptor antagonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with D3 (dopamine 3) receptor antagonist, are chosen from the group of diseases or disorders consisting of psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions,

malinger, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders, delirium and Parkinson disease.

5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive
10 behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D3 receptor antagonist
15 compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D3 receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

20 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D3 receptor antagonist compound to augment the
25 therapeutic effect or to provide a faster onset of the therapeutic effect of said D3 receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

30 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D3 receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D3 receptor antagonist compound,
35 further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D3 receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D3 receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said D3 receptor antagonist compound is chosen from the group consisting of BSF-201640 and PD 58491, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a D3 receptor antagonist, preferably chosen from the group consisting of BSF-201640 and PD 58491, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders, delirium and Parkinson disease.

30: combination therapy with a DA (dopamine) uptake inhibitor

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a DA (dopamine) uptake inhibitor, are chosen from the group of diseases or disorders consisting of substance related disorders and Parkinson disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying

emotion dysregulation of substance related disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a DA uptake inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said DA uptake inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a DA uptake inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said DA uptake inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said DA uptake inhibitor compound is chosen from the group consisting of safinamide and GBR 12909, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a D2 receptor antagonist, preferably chosen from the group consisting of safinamide and GBR 12909, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of substance related disorders and Parkinson disease.

31: combination therapy with an dopamine (receptor) agonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an dopamine (receptor) agonist, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders,

personality disorders, problems related to abuse or neglect, pain disorders and Parkinson disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders and problems related to abuse or neglect., characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a dopamine (receptor) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said dopamine (receptor) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a dopamine (receptor) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said dopamine (receptor) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a dopamine (receptor) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said dopamine (receptor) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said dopamine (receptor) agonist compound is chosen from the group consisting of sumanirole, SLV 308, sarizotan, S32504, rotigotine (preferably a Once-a-Day Transdermal Patch), ropinirole HCL (preferably controlled-release formulation), pramipexole, DAB452, cabergoline, bromocriptine, alaptide amantadine, bromocriptine, cabergoline lisuride and pergolide, preferably sumanirole, rotigotine (preferably a Once-a-Day Transdermal Patch), pergolide or ropinirole HCL (preferably controlled-release formulation), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, sumanirole is to be administered in a daily dose of between 4 and 16 mg of the active ingredient. Preferably, rotigotine (Once-a-Day Transdermal Patch) is to be administered in a daily dose of between 4.5 and 13.5 mg of the active ingredient. Preferably, ropinirole HCL (controlled-release formulation) is to be administered in a daily dose of between 0.75 and 24 mg of the active ingredient. Preferably, pergolide is to be administered in a daily dose of between 0.5 and 10 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a dopamine (receptor) agonist, preferably chosen from the group consisting of sumanirole, SLV 308, sarizotan, S32504, rotigotine (preferably a Once-a-Day Transdermal Patch), ropinirole HCL (preferably controlled-release formulation), pramipexole, DAB452, cabergoline, bromocriptine, alaptide amantadine, bromocriptine, cabergoline lisuride and pergolide, more preferably sumanirole, rotigotine (preferably a Once-a-Day Transdermal Patch), ropinirole HCL (preferably controlled-release formulation) or pergolide, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, problems related to abuse or neglect, pain disorders and Parkinson disease.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said dopamine (receptor) agonist is sumanirole, preferably provided in a unitary dose of between 4 and 16 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said dopamine (receptor) agonist is rotigotine (Once-a-Day Transdermal Patch), preferably provided in a unitary dose of between 4.5 and 13.5 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said dopamine (receptor) agonist is ropinirole HCL (controlled-release formulation), preferably provided in a unitary dose of between 0.75 and 24 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said dopamine (receptor) agonist is pergolide, preferably provided in a unitary dose of between 0.5 and 10 mg of the active ingredient.

32: combination therapy with a compound activating ERK (extracellular signal-related kinase)

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound that activates ERK (extracellular signal-related kinase), are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a

general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound that activates ERK (extracellular signal-related kinase) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound that activates ERK (extracellular signal-related kinase), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said compound that activates ERK (extracellular signal-related kinase) is CPI-1189 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, CPI-1189 is to be administered in a daily dose of between 50 and 100 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound that activates ERK (extracellular signal-related kinase), preferably CPI-1189 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said compound that activates ERK (extracellular signal-related kinase) is CPI-1189, preferably provided in a unitary dose of between 50 and 100 mg of the active ingredient.

33: combination therapy with a GABA (gamma-aminobutyric acid) agonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a

GABA (gamma-aminobutyric acid) agonist compound, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a GABA agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GABA agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said GABA agonist compound is nefiracetam or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a GABA agonist, preferably nefiracetam or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition,

substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

34: combination therapy with a GABA-A agonist compound

5 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a GABA-A (gamma-aminobutyric acid receptor A) agonist compound, are sleep disorders.

10 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of sleep disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a GABA-A agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GABA-A agonist
15 compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said GABA-A agonist compound is gaboxadol or a pro-drug or an active metabolite
20 thereof, or a pharmaceutically acceptable salt thereof. Preferably, gaboxadol is to be administered in a daily dose of between 5 and 20 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an GABA-A agonist, preferably gaboxadol or a pro-drug or an active metabolite thereof,
25 or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of sleep disorders.

The invention also relates to a pharmaceutical composition as described above wherein
30 pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said GABA-A agonist is Gaboxadol, preferably provided in a unitary dose of between 5 and 20 mg of the active ingredient.

35: combination therapy with a GABA-A modulator compound

35 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a

GABA-A (gamma-aminobutyric acid receptor A) modulator compound, are chosen from the group of diseases or disorders consisting of anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a GABA-A modulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GABA-A modulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said GABA-A modulator compound is chosen from the group consisting of zolpidem (preferably MR sustained-release version), zaleplon (preferably extended-release formulation), SL 65.1498, SEP174559, pagoclone, NGD 96-3, indiplon, eszopiclone, CP-730,330 (NGD 96-3) and ocinaplon, preferably zolpidem (preferably MR sustained-release version), zaleplon (preferably extended-release formulation), pagoclone, NGD 96-3, indiplon or eszopiclone, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, zolpidem MR sustained-release version is to be administered in a daily dose of between 10 and 20 mg of the active ingredient. Preferably, zaleplon extended-release is to be administered in a daily dose ranging between 2.5 and 20 mg of the active ingredient. Preferably, pagoclone is to be administered in a daily dose ranging between 7.5 and 60 mg of the active ingredient. Preferably, indiplon is to be administered in a daily dose of between 10 and 20 mg of the active ingredient. Preferably, eszopiclone is to be administered in a daily dose of between 2 and 3 mg of the active

ingredient. Preferably, ocinaplon is to be administered in a daily dose of between 10 and 60 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and
5 (b) a GABA-A modulator, preferably chosen from the group consisting of zolpidem (preferably MR sustained-release version), zaleplon (preferably extended-release formulation), SL 65.1498, SEP174559, pagoclone, NGD 96-3, indiplon, eszopiclone, CP-730,330 (NGD 96-3) and ocinaplon, preferably zolpidem (preferably MR sustained-release version), zaleplon (preferably extended-release formulation), pagoclone, NGD 96-3, indiplon
10 or eszopiclone, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders),
15 factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect.

The invention also relates to a pharmaceutical composition as described above wherein
20 pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said GABA-A modulator is zolpidem MR sustained-release version, preferably provided in a unitary dose of between 10 and 20 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein
25 pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said GABA-A modulator is zaleplon extended-release, preferably provided in a unitary dose of between 2.5 and 20 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein
30 pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said GABA-A modulator is Pagoclone, preferably provided in a unitary dose of between 7.5 and 60 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein
35 pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient

and wherein said GABA-A modulator is indiplon, preferably provided in a unitary dose of between 10 and 20 mg of the active ingredient.

5 The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said GABA-A modulator is eszopiclone, preferably provided in a unitary dose of between 2 and 3 mg of the active ingredient.

10 The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said GABA-A modulator is ocinaplon, preferably provided in a unitary dose of between 10 and 60 mg of the active ingredient.

36: combination therapy with a GABA-B antagonist compound

15 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a GABA-B (gamma-aminobutyric acid receptor B) antagonist compound, are chosen from the group of diseases or disorders consisting of anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious
20 disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect.

25 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders,
30 impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a GABA-B antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GABA-B antagonist
35 compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said GABA-B antagonist compound is AVE 7398 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a GABA-B antagonist, preferably AVE 7398 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect.

37: combination therapy with a Glial-cell Line Derived Neurotrophic Factor compound

The mental disorder which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a Glial-cell Line Derived Neurotrophic Factor compound, is Parkinson disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a Glial-cell Line Derived Neurotrophic Factor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said Glial-cell Line Derived Neurotrophic Factor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said Glial-cell Line Derived Neurotrophic Factor compound is GDNF or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, GDNF is to be administered in a daily dose ranging between 3.75 and 30 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a Glial-cell Line Derived Neurotrophic Factor, preferably GDNF or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson disease.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said Glial-cell Line Derived Neurotrophic Factor is GDNF, preferably provided in a unitary dose of between 3.75 and 30 mg of the active ingredient.

38: combination therapy with a glucocorticoid synthesis inhibitor compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a glucocorticoid synthesis inhibitor compound, are chosen from the group of diseases or disorders consisting of substance related disorders and Parkinson disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of substance related disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a glucocorticoid synthesis inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said glucocorticoid synthesis inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a glucocorticoid synthesis inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said glucocorticoid synthesis inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said glucocorticoid synthesis inhibitor compound is metyrapone or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

5 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a glucocorticoid synthesis inhibitor, preferably metyrapone or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion
10 dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of substance related disorders and Parkinson disease.

39: combination therapy with a glutamate receptor antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity
15 for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a glutamate receptor antagonist compound, are chosen from the group of diseases or disorders consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem,
20 problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group
25 of diseases and disorders consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously
30 with, separate from or prior to the administration of a glutamate receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said glutamate receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

35

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a glutamate receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said glutamate receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said glutamate receptor antagonist compound is LY354740 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a glutamate receptor antagonist, preferably LY354740 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

40: combination therapy with an GPCR (G-protein-coupled receptor) modulator

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an GPCR (G-protein-coupled receptor) modulator, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying

emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a GPCR modulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GPCR modulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a GPCR modulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GPCR modulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said GPCR modulator compound is R1204 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a GPCR modulator, preferably R1204 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

41: combination therapy with an GR (glucocorticoid receptor) antagonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an GR (glucocorticoid receptor) antagonist, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a GR antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GR antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a GR antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GR antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said GR antagonist compound is chosen from the group consisting of ORG 34517/34850 and mifepristone, preferably mifepristone, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, mifepristone is to be administered in a daily dose of between 600 and 1200 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a GR antagonist, preferably chosen from the group consisting of ORG 34517/34850 and mifepristone, preferably mifepristone, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said GR antagonist is Mifepristone, preferably provided in a unitary dose of between 600 and 1200 mg of the active ingredient.

42: combination therapy with a histamine H3-receptor antagonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a histamine H₃-receptor antagonist, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying

- emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a histamine H3-receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said histamine H3-receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.
- According to a preferred embodiment, the invention relates to the uses as described above, wherein said histamine H3-receptor antagonist compound is chosen from the group of compounds consisting of ABT-834 and ABT-239, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.
- The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a histamine H3-receptor antagonist, preferably chosen from the group consisting of ABT-834 and ABT-239 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is chosen from the group consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders.

43: combination therapy with a hormonal substance

- The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a hormonal substance, are chosen from the group of diseases or disorders consisting of premenstrual syndrome and sexual and gender identity disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of premenstrual syndrome and sexual and gender identity disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a hormonal substance to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said hormonal substance, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said hormonal substance is chosen from the group consisting of a testosterone transdermal spray, a testosterone gel, a female testosterone patch, synthetic conjugated estrogen A, methyltestosterone, a estrogens/methyltestosterone and a drosiperone/ethinyl estradiol composition, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said hormonal substance is synthetic conjugated estrogen A and is to be administered in a daily dose ranging between 0.075 and 0.6 mg of the active ingredient. More preferably, said hormonal substance is a drosiperone/ethinyl estradiol composition and is to be administered as a daily dose in tablets, preferably comprising 3mg drosiperone and 0.02 mg ethinyl estradiol of the active ingredients, respectively.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a hormonal substance, preferably chosen from the group consisting of a testosterone transdermal spray, a testosterone gel, a female testosterone patch, synthetic conjugated estrogen A, methyltestosterone, a estrogens/methyltestosterone and a drosiperone/ethinyl estradiol composition, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is chosen from the group consisting of premenstrual syndrome and sexual and gender identity disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient

and wherein said hormonal substance is synthetic conjugated estrogen A, preferably provided in a unitary dose of between 0.075 and 0.6 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein
5 pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said hormonal substance is a drosiperone/ethinyl estradiol composition, preferably provided in tablets comprising a unitary dose of 3mg drosiperone and 0.02 mg ethinyl estradiol of the active ingredients, respectively.

10 44: combination therapy with a compound which increases brain concentrations of 5-HT

The mental disorders which can be treated using compounds having a high selective affinity
15 for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound which increases brain concentrations of 5-HT (serotonin), are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating
15 disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

20 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders,
25 premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is
30 administered simultaneously with, separate from or prior to the administration of a compound which increases brain concentrations of 5-HT (serotonin) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which increases brain concentrations of 5-HT (serotonin), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active
35 ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which increases brain concentrations of 5-HT (serotonin) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which increases brain concentrations of 5-HT (serotonin), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said compound which increases brain concentrations of 5-HT (serotonin) is chosen from the group consisting of triptosine, tramadol, SP 186, PMD 145 and KW 6055, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which increases brain concentrations of 5-HT (serotonin), preferably chosen from the group consisting of triptosine, tramadol, SP 186, PMD 145 and KW 6055, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

45: combination therapy with a compound which increases insulin sensitivity

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound which increases insulin sensitivity, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical

condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which increases insulin sensitivity to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which increases insulin sensitivity, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said compound which increases insulin sensitivity is rosiglitazone maleate, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which increases insulin sensitivity, preferably rosiglitazone maleate or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is chosen from the group consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

46: combination therapy with a compound inhibiting the mixed lineage kinase family

The mental disorder which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound which is an inhibitor of the mixed lineage kinase family is Parkinson Disease.

5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from
10 or prior to the administration of a compound which is an inhibitor of the mixed lineage kinase family to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which is an inhibitor of the mixed lineage kinase family, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

15 According to a preferred embodiment, the invention relates to the use as described above, wherein said compound which is an inhibitor of the mixed lineage kinase family is CEP-1347 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

20 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which is an inhibitor of the mixed lineage kinase family, preferably CEP-1347 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson Disease.

25 47: combination therapy with an interleukin-1 beta converting enzyme inhibitor compound
The mental disorder which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an interleukin-1 beta converting enzyme inhibitor compound, is a pain disorder.

30 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a pain disorder, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from
35 or prior to the administration of an interleukin-1 beta converting enzyme inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said

interleukin-1 beta converting enzyme inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

- 5 According to a preferred embodiment, the invention relates to the use as described above, wherein said interleukin-1 beta converting enzyme inhibitor is pralnacasan or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

- 10 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an interleukin-1 beta converting enzyme inhibitor, preferably pralnacasan or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a pain disorder,.

15 48: combination therapy with a levodopa/decarboxylase inhibitor compound

The mental disorder which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a levodopa/decarboxylase inhibitor compound, is Parkinson Disease.

- 20 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a levodopa/decarboxylase inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said
- 25 levodopa/decarboxylase inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

- 30 According to a preferred embodiment, the invention relates to the use as described above, wherein said levodopa/decarboxylase inhibitor compound is levodopa/carbidopa, levodopa/benserazide, etilevodopa/carbidopa or etilevodopa/benserazide, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. According to a further preferred embodiment, the invention relates to the use as described above, wherein said
- 35 levodopa/decarboxylase inhibitor compound is (eti)levodopa/carbidopa, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof in combination with

entacapone, which is an inhibitor of catechol-O-methyltransferase (COMT), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably said levodopa/decarboxylase inhibitor compound is levodopa/carbidopa and is to be administered in a dose ranging between 2000 mg/ 50 mg and 100 mg/ 10 mg of the active ingredients.

5 Preferably said entacapone is to be administered in a dose ranging between 1000 mg/ 50 mg, more preferably between 500 mg/ 100 mg, and most preferably 200 mg of the active ingredients per day.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and

10 (b) a levodopa/decarboxylase inhibitor compound, preferably levodopa/carbidopa, levodopa/benserazide, etilevodopa/carbidopa or etilevodopa/benserazide, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson Disease. The invention also relates to a pharmaceutical

15 composition comprising (a) pipamperon, and (b) a levodopa/decarboxylase inhibitor compound, preferably is (eti)levodopa/carbidopa, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof in combination with entacapone, which is an inhibitor of catechol-O-methyltransferase (COMT), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for

20 simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson Disease.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said levodopa/decarboxylase inhibitor compound is levodopa/carbidopa,

25 preferably provided in a unitary dose of between 100 mg and 10 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said levodopa/decarboxylase inhibitor compound is levodopa/carbidopa or

30 etilevodopa/ carbidopa in combination with entacapone, of which the latter is preferably provided in a unitary dose of between 500 mg and 100 mg of the active ingredient.

49: combination therapy with a lipid-DNA complex

The mental disorder which can be treated using compounds having a high selective affinity

35 for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a lipid-DNA complex, is Parkinson Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of lipid-DNA complex to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said lipid-DNA complex, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said lipid-DNA complex is GR213487B or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a lipid-DNA complex, preferably GR213487B or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson Disease.

50: combination therapy with a monoamine oxidase (MAO) reuptake inhibitor

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a monoamine oxidase (MAO) reuptake inhibitor, are chosen from the group of diseases or disorders consisting of substance related disorders and attention-deficit disorders (ADHD).

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of non-cognitive mental disease or disorder which are substance related disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase (MAO) reuptake inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said monoamine oxidase (MAO) reuptake inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of attention-deficit disorders (ADHD), characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase (MAO) reuptake inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said monoamine oxidase (MAO) reuptake inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said monoamine oxidase (MAO) reuptake inhibitor compound is NS 2359 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a monoamine oxidase (MAO) reuptake inhibitor, preferably NS 2359 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of substance related disorders and attention-deficit disorders (ADHD).

51: combination therapy with a MAO-A and a MAO-B reuptake inhibitor

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a monoamine oxidase A (MAO-A) and a monoamine oxidase B (MAO-B) reuptake inhibitor, wherein said disorders are attention-deficit disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of attention-deficit disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase A (MAO-A) and a monoamine oxidase B (MAO-B) reuptake inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said monoamine oxidase A (MAO-A) and a monoamine oxidase B (MAO-B) reuptake inhibitor compound, further characterized in that

pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said monoamine oxidase A (MAO-A) and a monoamine oxidase B (MAO-B) reuptake inhibitor compound is SPD473 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a monoamine oxidase A (MAO-A) and a monoamine oxidase B (MAO-B) reuptake inhibitor, preferably SPD473 or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of attention-deficit disorders.

52: combination therapy with a monoamine oxidase B (MAO-B) inhibitor

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a monoamine oxidase B (MAO-B) inhibitor, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, problems related to abuse or neglect, pain disorder and Parkinson Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase B (MAO-B) inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said monoamine oxidase B (MAO-B) inhibitor compound, further characterized in that pipamperon is to be

administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase B (MAO-B) inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said monoamine oxidase B (MAO-B) inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase B (MAO-B) inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said monoamine oxidase B (MAO-B) inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said monoamine oxidase B (MAO-B) inhibitor compound is chosen from the group consisting of selegiline, rasagiline (TVP-1012) and EmSam (transdermal selegiline), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said monoamine oxidase B (MAO-B) inhibitor is selegiline and is to be administered in a daily dose ranging between 5 and 10 mg of the active ingredient. More preferably, said monoamine oxidase B (MAO-B) inhibitor is rasagiline (TVP-1012) and is to be administered in a daily dose ranging between 1 and 2 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a monoamine oxidase B (MAO-B) inhibitor, preferably chosen from the group consisting of selegiline, rasagiline (TVP-1012) and EmSam (transdermal selegiline), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined

D

preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, problems related to abuse or neglect, pain disorder and Parkinson Disease.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said monoamine oxidase B (MAO-B) inhibitor is selegiline, preferably provided in a unitary dose of between 5 and 10 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said monoamine oxidase B (MAO-B) inhibitor is rasagiline (TVP-1012), preferably provided in a unitary dose of between 1 and 2 mg of the active ingredient.

53: combination therapy with a monoamine oxidase B (MAO-B) reuptake inhibitor

The mental disorder which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a monoamine oxidase B (MAO-B) reuptake inhibitor, is Parkinson Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase B (MAO-B) reuptake inhibitor to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said monoamine oxidase B (MAO-B) reuptake inhibitor, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said monoamine oxidase B (MAO-B) reuptake inhibitor is safinamide or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a monoamine oxidase B (MAO-B) reuptake inhibitor, preferably safinamide or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson Disease.

54: combination therapy with a melanocortin-4 (MC4) receptor antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a melanocortin-4 (MC4) receptor antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a melanocortin-4 (MC4) receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said melanocortin-4 (MC4) receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said

pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a melanocortin-4 (MC4) receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said melanocortin-4 (MC4) receptor antagonist compound, further characterized in that
5 pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said melanocortin-4 (MC4) receptor antagonist compound is MCL0129, or a pro-
10 drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a melanocortin-4 (MC4) receptor antagonist compound, preferably MCL0129 or a pro-
15 drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders,
20 impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

55: combination therapy with a MCH receptor antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a
25 melanin concentrating hormone (MCH) receptor antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep
30 disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying
35 emotion dysregulation of a non-cognitive mental disease or disorder selected from the

group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a melanin concentrating hormone (MCH) receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said melanin concentrating hormone (MCH) receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a melanin concentrating hormone (MCH) receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said melanin concentrating hormone (MCH) receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said melanin concentrating hormone (MCH) receptor antagonist compound is SNAP-7941 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a melanin concentrating hormone (MCH) receptor antagonist compound, preferably SNAP-7941 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders,

adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

5 **56: combination therapy with a melatonin receptor (MT) agonist compound**

10 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a melatonin receptor (MT) agonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

15 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders),
20 factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a melatonin
25 receptor (MT) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said melatonin receptor (MT) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

30 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a melatonin receptor (MT) agonist compound to augment the
35 therapeutic effect or to provide a faster onset of the therapeutic effect of said melatonin receptor (MT) agonist compound, further characterized in that pipamperon is to be

administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said melatonin receptor (MT) agonist compound is chosen from the group consisting of ramelteon and agomelatine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said melatonin receptor (MT) agonist compound is agomelatine and is to be administered in a daily dose ranging between 25 and 50 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a melatonin receptor (MT) agonist compound, preferably ramelteon or agomelatine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said melatonin receptor (MT) agonist compound is agomelatine, preferably provided in a unitary dose of between 25 and 50 mg of the active ingredient.

57: combination therapy with a metabotropic glutamate receptor (MgluR) agonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a metabotropic glutamate receptor (MgluR) agonist compound, are chosen from the group of diseases or disorders consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a metabotropic glutamate receptor (MgluR) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said metabotropic glutamate receptor (MgluR) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a metabotropic glutamate receptor (MgluR) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said metabotropic glutamate receptor (MgluR) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said metabotropic glutamate receptor (MgluR) agonist compound is PRE703 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a metabotropic glutamate receptor (MgluR) agonist, preferably PRE703 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, substance-related

disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

58: combination therapy with a compound mimicking the effect of nerve growth factor (NGF)

5 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound which mimics the effect of nerve growth factor (NGF), are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, 10 dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease.

15 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, 20 dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from 25 or prior to the administration of a compound which mimics the effect of nerve growth factor (NGF) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which mimics the effect of nerve growth factor (NGF), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

30 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from 35 or prior to the administration of a compound which mimics the effect of nerve growth factor (NGF) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect

of said compound which mimics the effect of nerve growth factor (NGF), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5 According to a preferred embodiment, the invention relates to the uses as described above, wherein said compound which mimics the effect of nerve growth factor (NGF) is xaliproden or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said compound which mimics the effect of nerve growth factor (NGF) is xaliproden and is to be administered in a daily dose ranging between 1 and 2 mg of the
10 active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which mimics the effect of nerve growth factor (NGF), preferably xaliproden or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof,
15 as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease,
20 dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient
25 and wherein said compound which mimics the effect of nerve growth factor (NGF) is xaliproden, preferably provided in a unitary dose of between 1 and 2 mg of the active ingredient.

30 59: combination therapy with a muscarinic receptor partial agonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a muscarinic receptor partial agonist compound, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular
35 dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease,

dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, 10 dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from 15 or prior to the administration of a muscarinic receptor partial agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said muscarinic receptor partial agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

20 According to a preferred embodiment, the invention relates to the use as described above, wherein said muscarinic receptor partial agonist compound is sevimeline or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

25 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a muscarinic receptor partial agonist compound, preferably sevimeline or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of 30 Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and 35 other cognitive disorders.

60: combination therapy with a selective nor-adrenaline re-uptake inhibitor (NARI) compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a selective nor-adrenaline re-uptake inhibitor (NARI) compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, adjustment disorders, attention-deficit disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, adjustment disorders, attention-deficit disorders, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective nor-adrenaline receptor inhibitor (NARI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective nor-adrenaline re-uptake inhibitor (NARI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective nor-adrenaline re-uptake inhibitor (NARI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective nor-adrenaline re-uptake inhibitor (NARI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said selective nor-adrenaline re-uptake inhibitor (NARI) compound is chosen from the group consisting of reboxetine, atomoxetine hydrochloride, A 75200, 155U88, (S)-A 75200, tandamine, pirandamine, ciclazindol, fluparoxan, lortalamine, talsupram, talopram, prindamine, nomifensine, viloxazine, tomoxetine, duloxetine, venlafaxine and milnacipran or

a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said selective nor-adrenaline re-uptake inhibitor (NARI) compound is reboxetine and is to be administered in a daily dose ranging between 8 and 12 mg of the active ingredient. More preferably, said selective nor-adrenaline re-uptake inhibitor (NARI) compound is atomoxetine hydrochloride and is to be administered in a daily dose ranging between 40 and 100 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a selective nor-adrenaline re-uptake inhibitor (NARI) compound, preferably chosen from the group consisting of reboxetine, atomoxetine hydrochloride, A 75200, 155U88, (S)-A 75200, tandamine, pirandamine, ciclazindol, fluparoxan, lortalamine, talsupram, talopram, prindamine, nomifensine, viloxazine, tomoxetine, duloxetine, venlafaxine and milnacipran, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, adjustment disorders, attention-deficit disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective nor-adrenaline re-uptake inhibitor (NARI) compound is reboxetine, preferably provided in a unitary dose of between 8 and 12 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective nor-adrenaline re-uptake inhibitor (NARI) compound is atomoxetine hydrochloride, preferably provided in a unitary dose of between 40 and 100 mg of the active ingredient.

61: combination therapy with a NaSSA compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a noradrenergic/specific serotonergic antidepressant (NaSSA) compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating

disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders),
factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep
disorders, adjustment disorders, impulse control disorders, personality disorders, antisocial
behaviour, bereavement, occupational problem, problems related to abuse or neglect and
5 pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically
acceptable salt thereof for the preparation of a medicament for treating the underlying
emotion dysregulation of a non-cognitive mental disease or disorder selected from the
10 group of diseases and disorders consisting of mood disorders, anxiety disorders, eating
disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders),
factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep
disorders, adjustment disorders, impulse control disorders, personality disorders, antisocial
behaviour, bereavement, occupational problem and problems related to abuse or neglect,
15 characterized in that pipamperon or said pharmaceutically acceptable salt thereof is
administered simultaneously with, separate from or prior to the administration of a
noradrenergic/specific serotonergic antidepressant (NaSSA) compound to augment the
therapeutic effect or to provide a faster onset of the therapeutic effect of said
noradrenergic/specific serotonergic antidepressant (NaSSA) compound, further
20 characterized in that pipamperon is to be administered to a patient in a daily dose ranging
between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically
acceptable salt thereof for the preparation of a medicament for treating the underlying
25 emotion dysregulation of pain disorders, characterized in that pipamperon or said
pharmaceutically acceptable salt thereof is administered simultaneously with, separate from
or prior to the administration of a noradrenergic/specific serotonergic antidepressant
(NaSSA) compound to augment the therapeutic effect or to provide a faster onset of the
therapeutic effect of said noradrenergic/specific serotonergic antidepressant (NaSSA)
30 compound, further characterized in that pipamperon is to be administered to a patient in a
daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above,
wherein said noradrenergic/specific serotonergic antidepressant (NaSSA) compound is ORG
35 4420 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt
thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a noradrenergic/specific serotonergic antidepressant (NaSSA) compound, preferably ORG 4420 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

62: combination therapy with a selective NDRI compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, adjustment disorders, attention-deficit disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, adjustment disorders, attention-deficit disorders, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound to augment the therapeutic effect or to

provide a faster onset of the therapeutic effect of said selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

10

15

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

20

25

30

According to a preferred embodiment, the invention relates to the uses as described above, wherein said selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound is GW353162 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said selective nor-adrenaline and dopamine re-uptake inhibitor

35

(NDRI) compound is GW353162 and is to be administered in a daily dose ranging between 20 and 60 mg of the active ingredient.

5 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and
(b) a selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound,
preferably GW353162 or a pro-drug or an active metabolite thereof, or a pharmaceutically
acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential
10 use for treating the underlying emotion dysregulation of mental disease or disorder which is
chosen from the group consisting of mood disorders, anxiety disorders, adjustment
disorders, attention-deficit disorders, personality disorders, antisocial behaviour,
bereavement, occupational problem, problems related to abuse or neglect, pain disorders,
delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia,
15 dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson
Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due
to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition,
substance-induced persisting amnesic disorder, mild cognitive impairment disorder and
other cognitive disorders.

20 The invention also relates to a pharmaceutical composition as described above wherein
pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient
and wherein said selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI)
compound is GW353162, preferably provided in a unitary dose of between 20 and 60 mg of
the active ingredient.

25 63: combination therapy with a compound which is a neuroimmunophilin ligand

The mental disorder which can be treated using compounds having a high selective affinity
for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a
compound which is a neuroimmunophilin ligand, is Parkinson Disease.

30 The present invention thus relates to the use of pipamperon or a pharmaceutically
acceptable salt thereof for the preparation of a medicament for treating the underlying
emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said
pharmaceutically acceptable salt thereof is administered simultaneously with, separate from
or prior to the administration of a compound which is a neuroimmunophilin ligand to augment
35 the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound
which is a neuroimmunophilin ligand, further characterized in that pipamperon is to be

administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said a compound which is a neuroimmunophilin ligand is GPI 1485 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said a compound which is a neuroimmunophilin ligand is GPI 1485 and is to be administered in a daily dose ranging between 200 and 1000 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which is a neuroimmunophilin ligand, preferably GPI 1485 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson Disease.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said a compound which is a neuroimmunophilin ligand is GPI 1485, preferably provided in a unitary dose of between 200 and 1000 mg of the active ingredient.

64: combination therapy with a neuromodulator compound

The mental disorder which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a neuromodulator compound, is Parkinson Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neuromodulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neuromodulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said neuromodulator compound is adenosine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

- 5 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a neuromodulator compound, preferably adenosine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson Disease.

10 65: combination therapy with a neurotensin receptor antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a neurotensin receptor antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neurotensin receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurotensin receptor

antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neurotensin receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurotensin
10 receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

15 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neurotensin receptor antagonist compound to augment the therapeutic effect or to provide a
20 faster onset of the therapeutic effect of said neurotensin receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

25 According to a preferred embodiment, the invention relates to the use as described above, wherein said neurotensin receptor antagonist compound is SR 48692 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said neurotensin receptor antagonist compound is SR 48692 and is to be administered in a daily dose ranging between 90 and 300 mg of the active ingredient.

30 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a neurotensin receptor antagonist compound, preferably SR 48692 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting
35 mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment

disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

5

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said neurotensin receptor antagonist compound is SR 48692, preferably provided in a unitary dose of between 90 and 300 mg of the active ingredient.

10

66: combination therapy with nerve growth factor (NGF) gene therapy

15

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with nerve growth factor (NGF) gene therapy, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease.

20

25

30

35

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from nerve growth factor (NGF) gene therapy, to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said nerve growth factor (NGF) gene therapy, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to nerve growth factor (NGF) gene therapy, to augment the therapeutic effect or to provide a faster onset of nerve growth factor (NGF) gene therapy, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound useful in nerve growth factor (NGF) gene therapy, preferably xaliproden or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease.

It should be understood that "nerve growth factor gene therapy" is well known in the art, and the compounds, for instance nucleic acids used in nerve growth factor gene therapy are well described (see e.g. Tuszynski *et al.*, (2002) Journal of Molecular Neuroscience Volume 19, Issue 1-2, pps. 207-208).

67: combination therapy with a nicotinic acetylcholine receptor antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a nicotinic acetylcholine receptor antagonist compound, are chosen from the group of diseases or disorders consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a nicotinic acetylcholine receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said nicotinic acetylcholine receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a nicotinic acetylcholine receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said nicotinic acetylcholine receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said nicotinic acetylcholine receptor antagonist compound is SEP174559 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a nicotinic acetylcholine receptor antagonist compound, preferably SEP174559 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders,

substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

68: combination therapy with a nicotinic receptor agonist compound

- 5 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a nicotinic receptor agonist compound, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson
- 10 Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.
- 15 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma,
- 20 dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from
- 25 or prior to the administration of a nicotinic receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said nicotinic receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.
- 30 According to a preferred embodiment, the invention relates to the use as described above, wherein said nicotinic receptor agonist compound is ABT-089, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said nicotinic receptor agonist compound is ABT-089 and is to be administered in a daily dose ranging between 4 and 40 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a nicotinic receptor agonist compound, preferably ABT-089 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is chosen from the group consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The invention also relates to a pharmaceutical composition as described above, wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said nicotinic receptor agonist compound is ABT-089, preferably provided in a unitary dose of between 4 and 40 mg of the active ingredient.

69: combination therapy with a neurokinin 2 receptor (NK2) antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a neurokinin 2 receptor (NK2) antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is

administered simultaneously with, separate from or prior to the administration of a neurokinin 2 receptor (NK2) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurokinin 2 receptor (NK2) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neurokinin 2 receptor (NK2) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurokinin 2 receptor (NK2) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said neurokinin 2 receptor (NK2) antagonist compound is saredutant or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said neurokinin 2 receptor (NK2) antagonist compound is saredutant and is to be administered in a daily dose ranging between 25 and 200 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a neurokinin 2 receptor (NK2) antagonist compound, preferably saredutant or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient

and wherein said neurokinin 2 receptor (NK2) antagonist compound is saredutant, preferably provided in a unitary dose of between 25 and 200 mg of the active ingredient.

70: combination therapy with a neurokinin 3 receptor (NK3) antagonist compound

5 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a neurokinin 3 receptor (NK3) antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

15 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neurokinin 3 receptor (NK3) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurokinin 3 receptor (NK3) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

35 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from

or prior to the administration of a neurokinin 3 receptor (NK3) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurokinin 3 receptor (NK3) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neurokinin 3 receptor (NK3) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurokinin 3 receptor (NK3) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said neurokinin 3 receptor (NK3) antagonist compound is talnetant or osanetant, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said neurokinin 3 receptor (NK3) antagonist compound is talnetant and is to be administered in a daily dose ranging between 1.5 and 12 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a neurokinin 3 receptor (NK3) antagonist compound, preferably talnetant or osanetant, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said neurokinin 3 receptor (NK3) antagonist compound is talnetant, preferably provided in a unitary dose of between 1.5 and 12 mg of the active ingredient.

71: combination therapy with an N-Methyl-D-aspartate (NMDA) antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an N-Methyl-D-aspartate (NMDA) antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an N-Methyl-D-aspartate (NMDA) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said N-Methyl-D-aspartate (NMDA) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an N-Methyl-D-aspartate (NMDA) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said N-Methyl-D-aspartate (NMDA) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an N-Methyl-D-aspartate (NMDA) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said N-Methyl-D-aspartate (NMDA) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said N-Methyl-D-aspartate (NMDA) antagonist compound is chosen from the group consisting of SEP174559, memantine, delucemine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said N-Methyl-D-aspartate (NMDA) antagonist compound is memantine and is to be administered in a daily dose ranging between 5 and 40 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an N-Methyl-D-aspartate (NMDA) antagonist compound, preferably chosen from the group consisting of SEP174559, memantine, delucemine or a pro-drug or an active

metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said N-Methyl-D-aspartate (NMDA) antagonist compound is memantine, preferably provided in a unitary dose of between 5 and 40 mg of the active ingredient.

72: combination therapy with a non-steroidal anti-inflammatory drug

The mental disorder which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a non-steroidal anti-inflammatory drug, is a pain disorder or Alzheimer Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a pain disorder, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a non-steroidal anti-inflammatory drug to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said a non-steroidal anti-inflammatory drug, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive disease, such as Alzheimer Disease, characterized in

121

that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a non-steroidal anti-inflammatory drug to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said a non-steroidal anti-inflammatory drug, further characterized in that
5 pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said a non-steroidal anti-inflammatory drug is chosen from the group consisting of
10 piroxicam , MX-1094, meloxicam and flurizan (pure R-enantiomer form of flurbiprofen), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a non-steroidal anti-inflammatory drug, preferably chosen from the group consisting of
15 piroxicam, MX-1094, meloxicam and flurizan (pure R-enantiomer form of flurbiprofen), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a pain disorder or Alzheimer Disease .

20 73: combination therapy with an opoid antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an opoid antagonist compound, are substance related disorders.

25 It will be appreciated that the terms "opoid" and "opioid" may be used interchangeably.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of substance related disorders, characterized in that pipamperon or
30 said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a opoid antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said opoid antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

35

According to a preferred embodiment, the invention relates to the use as described above, wherein said opoid antagonist compound is naltrexone, preferably as a depot formulation, more preferably in the form of microcapsules, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, said naltrexone is to be administered in the form of a depot, preferably a depot of microcapsules comprising a daily dose of between 192 and 384 mg.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a opoid antagonist, preferably naltrexone, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of substance related disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said opoid antagonist compound is naltrexone, preferably provided in a unitary dose of between 192 and 384 mg of the active ingredient.

74: combination therapy with an opoid agonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an opoid agonist compound, are chosen from the group of diseases or disorders consisting of anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or

neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an opoid agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said opoid agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said opoid agonist compound is chosen from the group consisting of siramesine, E-5842 and cyclazocine, preferably siramesine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an opoid agonist compound, preferably chosen from the group consisting of siramesine, E-5842 and cyclazocine, preferably siramesine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder which is chosen from the group consisting of anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect.

75: combination therapy with a phosphodiesterase-4 (PDE4) inhibitor compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a phosphodiesterase-4 (PDE4) inhibitor compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general

medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of phosphodiesterase-4 (PDE4) inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said phosphodiesterase-4 (PDE4) inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

20 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a phosphodiesterase-4 (PDE4) inhibitor compound d to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said phosphodiesterase-4 (PDE4) inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

30 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive

125

impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a phosphodiesterase-4 (PDE4) inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said phosphodiesterase-4 (PDE4) inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said phosphodiesterase-4 (PDE4) inhibitor compound is chosen from the group consisting of ND1251 and MEM 1917 (R1497), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a phosphodiesterase-4 (PDE4) inhibitor antagonist compound, preferably chosen from the group consisting of ND1251 and MEM 1917 (R1497), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

76: combination therapy with a peptidic compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a peptidic compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and

gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a peptidic compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said peptidic compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a peptidic compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said peptidic compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting

dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a peptidic compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said peptidic compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said peptidic compound is chosen from the group consisting of secretin, PT-141, INN 00835 and beta sheet breaker peptide, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said peptidic compound is secretin and is to be administered in a daily dose ranging between 0.2 and 0.4 mg/kg of the active ingredient. More preferably, said peptidic compound is INN 00835 and is to be administered in a daily dose ranging between 18 and 160 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a peptidic compound, preferably chosen from the group consisting of secretin, PT-141, INN 00835 and beta sheet breaker peptide, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders.

128

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said peptidic compound is secretin, preferably provided in a unitary dose of 0.2 and 0.4 mg/kg of the active ingredient.

5

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said peptidic compound is INN 00835, preferably provided in a unitary dose of 18 and 160 mg of the active ingredient.

10

77: combination therapy with a phospholipase A2 inhibitor compound

15

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a phospholipase A2 inhibitor compound which has caspase inhibitor activity, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders and delirium.

20

25

30

35

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect,, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a phospholipase A2 inhibitor compound which has caspase inhibitor activity to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said phospholipase A2 inhibitor compound which has caspase inhibitor activity, further characterized in that

pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a phospholipase A2 inhibitor compound which has caspase inhibitor activity to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said phospholipase A2 inhibitor compound which has caspase inhibitor activity, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a phospholipase A2 inhibitor compound which has caspase inhibitor activity to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said phospholipase A2 inhibitor compound which has caspase inhibitor activity, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said phospholipase A2 inhibitor compound which has caspase inhibitor activity is chosen from the group consisting of LAX-101a, LAX-101b and LAX-101c, preferably LAX-101a, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a phospholipase A2 inhibitor compound which has caspase inhibitor activity, preferably chosen from the group consisting of LAX-101a, LAX-101b and LAX-101c, more preferably LAX-101a, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is

chosen from the group consisting mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders and delirium.

78: combination therapy with a compound which is a prodrug of uridine

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound which is a prodrug of uridine, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which is a prodrug of uridine to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which is a prodrug of uridine, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said

131

pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which is a prodrug of uridine to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which is a prodrug of uridine, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said compound which is a prodrug of uridine is RG2133 (triacetyluridine) or a prodrug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which is a prodrug of uridine, preferably RG2133 (triacetyluridine) or a prodrug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

79: combination therapy with prostaglandin E1 compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with prostaglandin E1 compound, are sexual and gender identity disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of sexual and gender identity disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a prostaglandin E1 compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said prostaglandin E1 compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

132

According to a preferred embodiment, the invention relates to the use as described above, wherein said prostaglandin E1 is alprostadil or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said prostaglandin E1 compound is alprostadil, preferably in the form of cream or gel, preferably a topical gel, and is to be administered in a daily dose ranging between 50 and 300 microgram per application of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a prostaglandin E1 compound, preferably alprostadil or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of sexual and gender identity disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said prostaglandin E1 compound is alprostadil, preferably provided in the form of a cream or gel, preferably a topical gel, wherein a unitary dose comprises between 50 and 300 microgram of the active ingredient per application.

80: combination therapy with a compound protecting dopaminergic and cholinergic neurons

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound which protects dopaminergic and cholinergic neurons, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma,

dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which protects dopaminergic and cholinergic neurons to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which protects dopaminergic and cholinergic neurons, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which protects dopaminergic and cholinergic neurons to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which protects dopaminergic and cholinergic neurons, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said compound which protects dopaminergic and cholinergic neurons is SR 57667 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which protects dopaminergic and cholinergic neurons, preferably SR 57667 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease.

81: combination therapy with a psychostimulant

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a psychostimulant, are chosen from the group of diseases or disorders consisting of sleep disorders, attention-deficit disorders and substance-related disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of sleep disorders, attention-deficit disorders and substance-related disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a psychostimulant to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said psychostimulant, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said psychostimulant is chosen from the group consisting of SPD 503, r-modafinil and modafinil, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said psychostimulant is SPE 503, more preferably said psychostimulant is modafinil and is to be administered in a daily dose ranging between 200 and 600 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a psychostimulant, preferably chosen from the group consisting of SPD 503, r-modafinil and modafinil, more preferably said SPC 503 or modafinil or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder which is chosen from the group consisting of sleep disorders, attention-deficit disorders and substance-related disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient

and wherein said psychostimulant is modafinil, preferably provided in a unitary dose of between 200 and 600 mg of the active ingredient.

82: combination therapy with a compound which is a Reversible Inhibitor of Mono-Amine oxydase A (RIMA)

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound which is a reversible inhibitor of mono-amine oxydase A (RIMA), are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which is a reversible inhibitor of mono-amine oxydase A (RIMA) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which is a reversible inhibitor of mono-amine oxydase A (RIMA), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which is a reversible Inhibitor of mono-amine

oxydase A (RIMA) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which is a reversible inhibitor of mono-amine oxydase A (RIMA), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5

According to a preferred embodiment, the invention relates to the uses as described above, wherein said compound which is a reversible inhibitor of mono-amine oxydase A (RIMA) is chosen from the group consisting of toloxatone, RS 8359, moclobemide, cimoxatone, caroxazone (F.I 6654) and befloxtatone, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said compound which is a reversible inhibitor of mono-amine oxydase A (RIMA) is befloxtatone and is to be administered in a daily dose ranging between 2.5 and 20 mg of the active ingredient.

10

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which is a reversible inhibitor of mono-amine oxydase A (RIMA), preferably chosen from the group consisting of toloxatone, RS 8359, moclobemide, cimoxatone, caroxazone (F.I 6654) and befloxtatone, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

15

20

25

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said compound which is a reversible inhibitor of mono-amine oxydase A (RIMA) is befloxtatone, preferably provided in a unitary dose of between 2.5 and 20 mg of the active ingredient.

30

83: combination therapy with a compound which modulates SCT-11

35

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound which modulates SCT-11 (i.e. SCT-11 is a G protein-coupled receptor), are

chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which modulates SCT-11 to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which modulates SCT-11, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which modulates SCT-11 to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which modulates SCT-11, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said compound which modulates SCT-11 is SNEC-2 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.







The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which modulates SCT-11, preferably SNE-2 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

84: combination therapy with a serotonin/dopamine antagonist compound (SDA)

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a serotonin/dopamine antagonist compound (SDA), are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a serotonin/dopamine antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said serotonin/dopamine

antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a serotonin/dopamine antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said serotonin/dopamine antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a serotonin/dopamine antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said serotonin/dopamine antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said serotonin/dopamine antagonist compound is chosen from the group consisting of zotepine, ziprasidone, SM-13496, SL 91.0177, sertindole, S-18327, risperidone, quetiapine fumarate (preferably sustained release formulation), quetiapine fumarate (preferably granules), quetiapine, perospirone, paliperidone, olanzapine, ocaperidone, LU 31-131, iloperidone, clozapine, BSF-190555, blonanserin, bifeprunox, asenapine and aripiprazole, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Even more preferably, said serotonin/dopamine antagonist compound is chosen from the group consisting of SL 91.0177, sertindole, perospirone, paliperidone, blonanserin, bifeprunox and asenapine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said serotonin/dopamine antagonist compound is sertindole and is to be administered in a daily dose ranging between 12 and 24 mg of the active ingredient. More preferably, said serotonin/dopamine antagonist

compound is paliperidone and is to be administered in a daily dose ranging between 3 and 15 mg of the active ingredient. More preferably, said serotonin/dopamine antagonist compound is asenapine and is to be administered in a daily dose ranging between 2.5 and 20 mg of the active ingredient.

5

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a serotonin/dopamine antagonist compound, preferably chosen from the group consisting of SL 91.0177, sertindole, perospirone, paliperidone, blonanserin, bifeprunox and asenapine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

10

15

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said serotonin/dopamine antagonist compound is sertindole, preferably provided in a unitary dose of between 12 and 24 mg of the active ingredient.

20

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said serotonin/dopamine antagonist compound is paliperidone, preferably provided in a unitary dose of between 3 and 15 mg of the active ingredient.

25

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said serotonin/dopamine antagonist compound is asenapine, preferably provided in a unitary dose of between 2.5 and 20 mg of the active ingredient.

30

85: combination therapy with a selective SDRI compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a selective serotonin and dopamine re-uptake inhibitor (SDRI) compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin and dopamine reuptakeinhibitor (SDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin and dopamine reuptake inhibitor (SDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from

or prior to the administration of a selective serotonin and dopamine reuptake inhibitor (SDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin and dopamine reuptake inhibitor (SDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin and dopamine reuptake inhibitor (SDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin and dopamine reuptake inhibitor (SDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said selective serotonin and dopamine reuptake inhibitor (SDRI) compound is bazinaprine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a selective serotonin and dopamine reuptake inhibitor (SDRI) compound, preferably bazinaprine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, antisocial behaviour,

bereavement, occupational problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

86: combination therapy with a second messenger beta agonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a second messenger beta agonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a second messenger beta agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said second messenger beta agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said

pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a second messenger beta agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said second messenger beta agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said second messenger beta agonist compound is chosen from the group consisting of SR 57227, rolipram and eplivanserin, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said second messenger beta agonist compound is rolipram and is to be administered in a daily dose ranging between 1.5 and 3 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a second messenger beta agonist compound, preferably chosen from the group consisting of SR 57227, rolipram and eplivanserin or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said second messenger beta agonist compound is rolipram, preferably provided in a unitary dose of between 1.5 and 3 mg of the active ingredient.

87: combination therapy with a secretin pancreatic hormone

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a secretin pancreatic hormone, are chosen from the group of diseases or disorders consisting of anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders),

factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a secretin pancreatic hormone to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said secretin pancreatic hormone, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a secretin pancreatic hormone to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said secretin pancreatic hormone, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a secretin

pancreatic hormone to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said secretin pancreatic hormone, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5

According to a preferred embodiment, the invention relates to the uses as described above, wherein said secretin pancreatic hormone is RG1068 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

- 10 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a secretin pancreatic hormone, preferably RG1068, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

20

88: combination therapy with a sigma receptor agonist compound

- The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a sigma receptor agonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

30

- The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders),

35

factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a sigma receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said sigma receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a sigma receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said sigma receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said sigma receptor agonist compound is VPI-013 (also known as OPC-14523) or PRX-00023, preferably VPI-013 (also known as OPC-14523), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a sigma receptor agonist compound, preferably VPI-013 (also known as OPC-14523) or PRX-00023, preferably VPI-013 (also known as OPC-14523), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

89: combination therapy with a sigma receptor antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a sigma receptor antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders and delirium.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect,, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a sigma receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said sigma receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a sigma receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said sigma receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a sigma receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said sigma receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said sigma receptor antagonist compound is chosen from the group consisting of SR 31742 and EMD 68843, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said sigma receptor antagonist compound is EMD 68843 and is to be administered in a daily dose ranging between 5 and 40 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a sigma receptor antagonist compound, preferably chosen from the group consisting of SR 31742 and EMD 68843, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders and delirium.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said sigma receptor antagonist compound is EMD 68843, preferably provided in a unitary dose of between 5 and 40 mg of the active ingredient.

90: combination therapy with a selective SNDRI compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from

or prior to the administration of a selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound is selected from the group consisting of NS 2330; McN 5652; DOV 216,303 and DOV 21,947; more preferably NS 2330 or DOV 216,303; or a pro-drug or an active metabolite thereof; or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound, preferably selected from the group consisting of NS 2330; McN 5652; DOV 216,303 and DOV 21,947, more preferably NS 2330 or DOV 216,303, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion

dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders.

91: combination therapy with a selective SNRI compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective

serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

15 According to a preferred embodiment, the invention relates to the uses as described above, wherein said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is selected from the group consisting of venlafaxine, tomoxetine, tandamine, talsupram, talopram, nefazodone, milnacipran, LY 113.821, duloxetine, desvenlafaxine and amoxapine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

20 Even more preferably, said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is chosen from the group consisting of venlafaxine, tomoxetine, milnacipran, duloxetine and desvenlafaxine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is venlafaxine and is to be administered in a daily dose ranging between 75 and 300 mg of the active ingredient. More preferably, said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is tomoxetine and is to be administered in a daily dose ranging between 0.475 and 3.8 mg/kg of the active ingredient. More preferably, said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is milnacipran and is to be administered in a daily dose ranging between

25 50 and 200 mg of the active ingredient. More preferably, said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is duloxetine and is to be administered in a daily dose ranging between 40 and 60 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and

35 (b) a selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound, preferably selected from the group consisting of venlafaxine, tomoxetine, tandamine, talsupram,

talopram, nefazodone, milnacipran, LY 113.821, duloxetine, desvenlafaxine and amoxapine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is venlafaxine, preferably provided in a unitary dose of between 75 and 300 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is tomoxetine, preferably provided in a unitary dose of between 0.475 and 3.8 mg/kg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is milnacipran, preferably provided in a unitary dose of between 50 and 200 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is duloxetine, preferably provided in a unitary dose of between 40 and 60 mg of the active ingredient.

92: combination therapy with a selective serotonin re-uptake inhibitor (SSRI) compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a selective serotonin re-uptake inhibitor (SSRI) compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin re-uptake inhibitor (SSRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin re-uptake inhibitor (SSRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin re-uptake inhibitor (SSRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin re-uptake inhibitor (SSRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said selective serotonin re-uptake inhibitor (SSRI) compound is selected from the group consisting of YM 992, VPI-013 (also known as OPC-14523), sertraline, paroxetine, LY 214.281, LU AA 21-004, Lu 35-138, litoxetine, ifoxetine, fluvoxamine (controlled release formulation), fluvoxamine, fluoxetine, femoxetine, escitalopram, EMD 68843, cyanodothepine, citalopram, venlafaxine, milnacipran, duloxetine, cericlamine and ademethionine (preferably s-adenosylmethionine), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Even more preferably, said selective serotonin re-uptake inhibitor (SSRI) compound is chosen from the group consisting of litoxetine, fluvoxamine (controlled release formulation) and escitalopram, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said selective serotonin re-uptake inhibitor (SSRI) compound is fluvoxamine (controlled release formulation) and is to be administered in a daily dose ranging between 100 and 300 mg of the active ingredient. More preferably, said selective serotonin re-uptake inhibitor (SSRI) compound is escitalopram and is to be administered in a daily dose ranging between 10 and 20 mg of the active ingredient. More preferably, said selective serotonin re-uptake inhibitor (SSRI) compound is citalopram and is to be administered in a daily dose ranging between 10 and 40 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a selective serotonin re-uptake inhibitor (SSRI) compound, preferably selected from the group consisting of YM 992, VPI-013 (also known as OPC-14523), sertraline, paroxetine, LY 214.281, LU AA 21-004, Lu 35-138, litoxetine, ifoxetine, fluvoxamine (controlled release formulation), fluvoxamine, fluoxetine, femoxetine, escitalopram, EMD 68843, cyanodothepine, citalopram, venlafaxine, milnacipran, duloxetine, cericlamine and ademethionine (preferably s-adenosylmethionine), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

157

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective serotonin re-uptake inhibitor (SSRI) compound is fluvoxamine (controlled release formulation), preferably provided in a unitary dose of between 100 and 300 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective serotonin re-uptake inhibitor (SSRI) compound is escitalopram, preferably provided in a unitary dose of between 10 and 20 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective serotonin re-uptake inhibitor (SSRI) compound is citalopram, preferably provided in a unitary dose of between 10 and 40 mg of the active ingredient.

Citalopram or citalopram hydrobromide is a selective serotonin (5-hydroxytryptamine / 5-HT) re-uptake inhibitor (SSRI) and is the conventional name given for the compound of the formula (RS)-1-[3-(dimethylamino)propyl]-1-(p-fluorophenyl)-5-phthalanecarbonitrile-hydrobromide. According to an embodiment, a daily doses of active ingredient of SSRI, preferably citalopram, ranges between 10 and 40 mg per day. Preferably, daily doses of active ingredient ranging between 20 and 30 mg per day are administered. More preferably, a daily dose of 10, 15, 20, 25, 30, 35 or 40 mg per day is administered.

Fluvoxamine or fluvoxamine maleate (luvox, fevarin) is a selective serotonin (5-HT) re-uptake inhibitor (SSRI) belonging to a new chemical series, the 2-aminoethyl oxime ethers of aralkylketones. It is chemically unrelated to other SSRIs and clomipramine. It is chemically designated as 5-methoxy-4'-(trifluoromethyl) valerophenone (E)-O-(2-aminoethyl) oxime maleate (1:1).

According to an embodiment, a daily dose of active ingredient of fluvoxamine in a controlled release mode ranges between 100 and 300 mg per day. Preferably, daily doses of active ingredient ranging between 150 and 200 mg per day are administered in a controlled release mode. More preferably, a daily dose of 100, 150, 200, 250 or 300 mg per day is administered by controlled release.

93: combination therapy with a substance P receptor (NK1) antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a substance P receptor (NK1) antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a substance P receptor (NK1) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said substance P receptor (NK1) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a substance P receptor (NK1) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said substance P receptor (NK1) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said substance P receptor (NK1) antagonist compound is chosen from the group consisting of vestipitant, TAK-637, R673, GW823296, GW679769, GW597599, CP-122.721, aprepitant, 823296 and 679769, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said substance P receptor (NK1) antagonist compound is aprepitant and is to be administered in a daily dose ranging between 40 and 160 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a substance P receptor (NK1) antagonist compound, preferably chosen from the group consisting of vestipitant, TAK-637, R673, GW823296, GW679769, GW597599, CP-122.721, aprepitant, 823296 and 679769, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said substance P receptor (NK1) antagonist compound is aprepitant, preferably provided in a unitary dose of between 40 and 160 mg of the active ingredient.

94: combination therapy with a sulfonamide compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a sulfonamide compound, are chosen from the group of diseases or disorders consisting of mood disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a sulfonamide compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said sulfonamide compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a sulfonamide compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said sulfonamide compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a sulfonamide compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said sulfonamide compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said sulfonamide compound is zonisamide or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said sulfonamide compound is zonisamide and is to be administered in a daily dose ranging between 100 and 600 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a sulfonamide compound, preferably zonisamide, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting mood disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said sulfonamide compound is zonisamide, preferably provided in a unitary dose of between 100 and 600 mg of the active ingredient.

95: combination therapy with a tachykinin antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a tachykinin antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group

D

of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a tachykinin antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said tachykinin antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a tachykinin antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said tachykinin antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said tachykinin antagonist compound is SR 48968 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a tachykinin antagonist compound, preferably SR 48968 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

96: combination therapy with a compound selected from the group consisting of R228060 (YKP-10A), palanpanel, ORG 39479/PH80, ORG 34167, DP 543 and CJ-017.493

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound selected from the group consisting of R228060 (YKP-10A), palanpanel, ORG 39479/PH80, ORG 34167, DP 543 and CJ-017.493, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, attention-deficit disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, phase of life problem, academic problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, attention-deficit disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, phase of life problem, academic problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound selected from the group consisting of R228060 (YKP-10A), palanpanel, ORG 39479/PH80, ORG 34167, DP 543 and CJ-017.493, to augment the therapeutic effect or to provide a

5 faster onset of the therapeutic effect of said compound selected from the group consisting of R228060 (YKP-10A), palanpanel, ORG 39479/PH80, ORG 34167, DP 543 and CJ-017.493, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound selected from the group consisting of R228060 (YKP-10A), palanpanel, ORG 39479/PH80, ORG 34167, DP 543 and CJ-017.493, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation
10 for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders,
15 impulse control disorders, pervasive development disorders, attention-deficit disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, phase of life problem, academic problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease,
20 substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders
25 and Parkinson Disease.

97: combination therapy with a vasopressin 1B receptor (V1B) antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a
30 vasopressin 1B receptor (V1B) antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality
35 disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a vasopressin 1B receptor (V1B) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said vasopressin 1B receptor (V1B) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a vasopressin 1B receptor (V1B) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said vasopressin 1B receptor (V1B) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said vasopressin 1B receptor (V1B) antagonist compound is SSR149415 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a vasopressin 1B receptor (V1B) antagonist compound, preferably SSR149415 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual

syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

5

98: combination therapy with a voltage-gated calcium channel $\alpha(2)\delta$ subunit modulator compound

10 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a voltage-gated calcium channel $\alpha(2)\delta$ subunit modulator compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, 15 personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

20 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, 25 characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a voltage-gated calcium channel $\alpha(2)\delta$ subunit modulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said voltage-gated calcium channel $\alpha(2)\delta$ subunit modulator compound, further characterized in that 30 pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

35 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said

pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a voltage-gated calcium channel alpha(2)delta subunit modulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said voltage-gated calcium channel alpha(2)delta subunit modulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said voltage-gated calcium channel alpha(2)delta subunit modulator compound is pregabalin or PD-200,390; or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said voltage-gated calcium channel alpha(2)delta subunit modulator compound is pregabalin, and is to be administered in a daily dose ranging between 50 and 600 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a voltage-gated calcium channel alpha(2)delta subunit modulator compound, preferably pregabalin or PD-200,390; or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said voltage-gated calcium channel alpha(2)delta subunit modulator compound is pregabalin, preferably provided in a unitary dose of between 50 and 600 mg of the active ingredient.

99: combination therapy with a vomeropherin compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a vomeropherin compound, are chosen from the group of diseases or disorders consisting of

anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of vomeropherin compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said vomeropherin compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said vomeropherin compound is PH94B or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) vomeropherin compound, preferably PH94B or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder which is chosen from the group consisting of anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect.

Also, the invention relates in particular to the use as described before, wherein said second compound is chosen from the group consisting of fluvoxamine controlled release,

phenserine tartrate, atomoxetine hydrochloride, bupropion (controlled-release formulation), ropinirole HCL (controlled-release formulation), INN 00835, galantamine (extended release formulation), paliperidone, tomoxetine, aprepitant, rivastigmine tartrate, ORG 34517/34850, sunepitron, sumanirole, milnacipran, idazoxan, xaliproden, SR 58611, befloxatone, litoxetine, 5 tianeptine, agomelatine, SPD 503, flesinoxan, bifeprunox, ramelteon, etilevodopa, rasagiline (TVP-1012) and desvenlafaxine.

Also, the invention relates in particular to the use as described before, wherein said second compound is chosen from the group consisting of galantamine (extended release formulation), R121919, risperidone, paliperidone and R228060 (YKP-10A).

- 10 The disclosure of all patents, publications (including published patent publications), and database accession numbers and depository accession numbers referenced in this specification are specifically incorporated herein by reference in their entirety to the same extent as if each such individual patent, publication, and database accession number, and depository accession number were specifically and individually indicated to be incorporated 15 by reference.

The invention, now being generally described, will be more readily understood by reference to the following tables and examples, which are included merely for purposes of illustration of certain aspects and embodiments of the present invention and are not intended to limit the invention.

Short description of the Tables and Figures

Table 1: In Table 1, the pKi values of test compounds are given for each of the dopamine receptors, 5HT receptors, adrenergic receptors and the histamine1 receptor.

5

Table 2: Set-up of a clinical trial comprising for treatment groups.

Table 3: Overview of a placebo, active and period controlled clinical trial in a fore-going pipamperon - citalopram treatment in Major Depressive Disorder.

10

Table 4: POC process for major depressive disorder.

Table 5: Summary of diseases and disorders relative to known psycho-tropics.

15

Table 6: Overview of Pharmacological grouping, indicating pharmacological profile numbering (column 2), pharmacological profile (column 3), main indication(s) (column 4), name of the compound (column 4), the dose range (column 5), and the company producing or selling said compound (column 6). Compounds indicated by hatching are preferred.

20

Figure 1: Add-on treatment with pipamperon after treatment with citalopram .

Figure 2: HDRS-17 change from baseline: combo treatment pipamperon as add-on - citalopram vs SNRI (duloxetine) in Major Depression.

25

Figure 3: Remission rates (HDRS-17 ≤ 7): combo treatment pipamperon as add-on - citalopram vs SNRI (venlafaxine) vs SSRIs vs placebo in Major Depression.

Figure 4: Fore-going treatment during 1-5 days with pipamperon followed with the combination treatment of pipamperon and citalopram.

30

Figure 5: HDRS-17 change from baseline: combo treatment pipamperon - citalopram with a fore-going treatment of 4 days with pipamperon vs SNRI (duloxetine) in Major Depression.

35

Figure 6: Remission rates (HDRS-17 ≤ 7): combo pipamperon - citalopram with a fore-going treatment of 4 days with pipamperon vs SNRI (venlafaxine) in Major Depression.

Figure 7: Fore-going treatment during 6-8 days with pipamperon followed with the combination treatment of pipamperon and citalopram.

Figure 8: HDRS-17 change from baseline: combo treatment pipamperon - citalopram with a fore-going treatment of 7 days with pipamperon vs SNRI (duloxetine) in Major Depression.

Figure 9: Fore-going and add-on treatment with pipamperon in MDD.

Figure 10: HDRS-17 change from baseline: fore-going and add-on treatment with pipamperon and citalopram in comparison with the SNRI duloxetine in Major Depression.

Figure 11: Remission rates (HDRS-17 ≤ 7): fore-going and add-on treatment with pipamperon and citalopram in comparison with the SNRI venlafaxine in Major Depression.

Figure 12: Y-BOCS total score: fore-going and add-on treatment with pipamperon and citalopram in comparison with the SSRI fluvoxamine in OCD.

Figure 13: Y-BOCS obsession score: fore-going and add-on treatment with pipamperon and citalopram in comparison with the SSRI fluvoxamine in OCD.

Figure 14: Y-BOCS compulsion score: fore-going and add-on treatment with pipamperon and citalopram in comparison with the SSRI fluvoxamine in OCD.

Table 1

Figure 15: CGI-severity score:	D1	D2	D3	D4	SHT _{1A}	SHT _{1B}	SHT _{1D}	SHT _{1E}	SHT _{1F}	SHT _{2A}	SHT _{2B}	SHT _{2C}	SHT _{6rat}	SHT _{7rat}	Alpha1	Alpha2	Alpha2	Alpha2	Beta1	Beta2	H1
ORG5222	8-9	8-9	8-9	8-9	8-9	8-9	8-9	7-8	0	<9	<9	<9	<9	<9	<9	8-9	8-9	7-8	<9	<9	<9
Zotepine	0	8-9	8-9	8-9	6-7	7-8	7-8	6-7	0	8-9	0	0	0	0	0	6-7	8-9	6-7	<9	<9	<9
Fluparoxan	0	<6	<6	<6	6-7	<6	<6	0	0	8-9	0	<6	0	0	6-7	8-9	8-9	8-9	0	0	0
Olazapine	7-8	7-8	7-8	7-8	<6	6-7	6-7	<6	6-7	8-9	8-9	8-9	7-8	6-7	7-8	6-7	6-7	6-7	<9	<9	<9
Clozapine	7-8	6-7	6-7	6-7	6-7	6-7	6-7	6-7	6-7	8-9	8-9	7-8	7-8	7-8	8-9	7-8	7-8	7-8	<6	<6	<9
S16924	0	7-8	7-8	7-8	8-9	0	0	0	0	>9	8-9	7-8	7-8	7-8	8-9	6-7	7-8	6-7	<6	<6	0
S18327	7-8	7-8	6-7	8-9	7-8	0	0	0	0	8-9	0	6-7	0	0	>9	6-7	0	0	0	0	0
Amperozide	6-7	6-7	6-7	8-9	<6	0	0	0	0	8-9	0	<6	0	0	7-8	<6	0	0	0	0	0
GGR218231	<6	7-8	>9	8-9	6-7	<6	<6	0	0	8-9	<6	<6	0	0	<6	6-7	6-7	6-7	<6	<6	6-7
Serindole	7-8	8-9	8-9	8-9	6-7	7-8	7-8	6-7	6-7	>9	0	8-9	0	0	>9	6-7	0	0	0	0	0
MDL100,907	6-7	<6	<6	8-9	<6	0	0	0	0	>9	0	7-8	0	0	<6	6-7	6-7	<6	<6	<6	6-7
Haloperidol	8-9	>9	8-9	8-9	<6	6-7	6-7	<6	<6	8-9	<6	<6	<6	6-7	8-9	<6	6-7	<6	<6	<6	6-7
Tiospirone	7-8	8-9	8-9	8-9	8-9	0	0	0	0	>9	0	8-9	0	0	>9	6-7	0	0	0	0	0
Raclopride	<6	8-9	8-9	8-9	<6	0	0	0	0	8-9	0	<6	0	0	<6	6-7	0	0	0	0	0
Fluspirilene	0	>9	8-9	8-9	7-8	<6	<6	<6	0	8-9	0	0	0	0	0	6-7	7-8	7-8	6-7	6-7	7-8
Ocaperidone	7-8	>9	8-9	8-9	7-8	0	0	0	0	>9	0	7-8	0	0	>9	0	0	0	0	0	0
Risperidone	7-8	8-9	7-8	8-9	6-7	8-9	6-7	<6	<6	>9	0	7-8	0	0	>9	7-8	8-9	8-9	<6	<6	7-8
S33084	6-7	7-8	>9	8-9	<6	6-7	6-7	0	0	8-9	6-7	7-8	0	0	6-7	<6	0	0	0	0	0
L741626	6-7	8-9	7-8	8-9	<6	<6	<6	<6	0	8-9	6-7	<6	0	0	6-7	<6	0	0	0	0	0
Seroquel	6-7	6-7	6-7	8-9	6-7	<6	<6	<6	<6	8-9	6-7	6-7	0	6-7	7-8	<6	7-8	6-7	<6	<6	8-9
Yohimbine	0	6-7	<6	8-9	7-8	6-7	6-7	0	0	8-9	0	<6	0	0	6-7	8-9	8-9	>9	<6	<6	0
Ziprasidone	8-9	8-9	7-8	7-8	8-9	>9	8-9	6-7	0	>9	8-9	8-9	7-8	8-9	8-9	6-7	7-8	7-8	<6	<6	7-8
Pipamperon	0	6-7	6-7	8-9	<6	6-7	6-7	>9	>9	8-9	0	0	0	0	0	6-7	7-8	7-8	<6	<6	<6

Table 2

173

ACUTE PHASE**										EXTENSION PHASE***										FOLLOW-UP PHASE			
VISITS	V1 Screen minus D7	V2 Baseline D0	V3 D4	V4 D7	V5 W2	V6 W3	V7 W4	V8 W6	V9 W8	V10 W10	V11 W12	V12 W16	V13 W20	V14 W24	V15 M8	V16 M10	V17 M12	V18 W1	V19 W2				
TREATMENT GROUP																							
Group Pip-Active / D7	A	B	B	C														A	A				
Group Pip-Active / D4	A	B	C															A	A				
Group Pip-Active / D0	A	C																A	A				
Group Plc-Active / D0	A	D																A	A				
Informant Consent	X																						
Weight	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X				
Physician Screen	X																						
CGI-S***	X																						
Q-LES-Q****	X																						
TOPIRAMATE	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X				

Treatment regimen:

A: PLC + PLC

B: 2x(PLC+PIP(4mg))/d

C: 2x(CIT(10mg)+PIP(4mg))/d

D: 2x(CIT(10mg)+PLC)/d

*Neuronal E-Clinical Trial = Vesalius Expert Development for this Trial which includes the bottom-up measurement of:

**Entering Acute Phase: only NON-placebo responders as defined by the DSM-IV criteria of efficacy

***Entering Extension Phase: only remitters as defined by the DSM-IV criteria of efficacy

****CGI-S : Clinical Global Impressions-Improvement Scale

*****Q-LES-Q : Quality of Life, Enjoyment and Satisfaction Questionnaire

Table 3

FOREGOING PIPAMPERON - CITALOPRAM TREATMENT IN MAJOR DEPRESSIVE DISORDER :																					
A PLACEBO, ACTIVE AND PERIODICALLY CONTROLLED																					
CLINICAL TRIAL																					
FOLLOW-UP PHASE																					
EXTENSION PHASE																					
ACUTE PHASE																					
VISITS		V1	V2	V3	V4	V5	V6	V7	V8	V9	V10	V11	V12	V13	V14	V15	V16	V17	V18	V19	
DAY / WEEK / MONTH		SCREEN minus D7	BASELINE D0	D4	D7	W2	W3	W4	W5	W6	W7	W8	W9	W10	W11	W12	W16	W20	M8	M10	M12
TREATMENT GROUP																					
Group PIP - Active / Day 4		PLC+PLC	2x(PLC+PIP(4mg))d	2x(CIT(10mg)+PIP(4mg))d																PLC+PLC	PLC+PLC
Group PIP - Active / Day 0		PLC+PLC	2x(CIT(10mg)+PIP(4mg))d																	PLC+PLC	PLC+PLC
Group PIP - Active / Day 0		PLC+PLC	2x(CIT(10mg)+PLC)d																	PLC+PLC	PLC+PLC
Group Placebo		PLC+PLC																			
Informed Consent		X																			
NEGOT		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Vital Signs / Weight		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
LAB		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
ECG		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Phys Exam		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Alc/Drugs Screen		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
CGIS		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
***Q-LES-Q		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X

*Neuronal E-Clinical Trial = Vesalius Expert Development for this Trial which includes the bottom-up measurement of:

- . In- and exclusion criteria
- . Functional status evaluation
- . Medical history
- . (Pre-)treatment signs & symptoms
- . DSM-IV rules for diagnosis & efficacy
- . Rating Scales: HDRS-20, MADRS, HAMA
- . Medical resource utilisation
- . Pre-trial & concomitant medication
- . Drug administration
- . (Serious) Adverse events
- . Admission to the acute and extension phase of treatment
- . Right flow of the trial

**Entering Acute Phase: only NON-placebo responders as defined by the DSM-IV criteria of efficacy

***Entering Extension Phase: only remitters as defined by the DSM-IV criteria of efficacy

****Q-LES-Q: Quality of Life, Enjoyment and Satisfaction Questionnaire

Table 4

DAY	minus D7	D0	=>D4
TREATMENTGROUP			
Placebo (PLC)	PLC+PLC	2x(PLC+PLC)	2x(PLC+PC)
PIP - Active / Day 4	PLC+PLC	2x(PLC+PIP (4mg))/d	2x(CIT (10mg)+PIP (4mg))/d
PIP - Active / Day 0	PLC+PLC	2x(CIT (10mg)+PIP (4mg))/d	2x(CIT (10mg)+PIP (4mg))/d
PLC - Active / Day 0	PLC+PLC	2x(CIT (10mg)+PLC)/d	2x(CIT (10mg)+PLC)/d

[illegible]

[illegible]

MEDICATION										13	15	17	19	21	23	25	27	29	31	33	35	37	39	41	43	45	47	49	51	53	55	57	59	61	63	65	67	69	71	73	75	77	79	81	83	85	87	89	91	93	95	97	99	101	103	105	107	109	111	113	115	117	119	121	123	125	127	129	131	133	135	137	139	141	143	145	147	149	151	153	155	157	159	161	163	165	167	169	171	173	175	177	179	181	183	185	187	189	191	193	195	197	199	201	203	205	207	209	211	213	215	217	219	221	223	225	227	229	231	233	235	237	239	241	243	245	247	249	251	253	255	257	259	261	263	265	267	269	271	273	275	277	279	281	283	285	287	289	291	293	295	297	299	301	303	305	307	309	311	313	315	317	319	321	323	325	327	329	331	333	335	337	339	341	343	345	347	349	351	353	355	357	359	361	363	365	367	369	371	373	375	377	379	381	383	385	387	389	391	393	395	397	399	401	403	405	407	409	411	413	415	417	419	421	423	425	427	429	431	433	435	437	439	441	443	445	447	449	451	453	455	457	459	461	463	465	467	469	471	473	475	477	479	481	483	485	487	489	491	493	495	497	499	501	503	505	507	509	511	513	515	517	519	521	523	525	527	529	531	533	535	537	539	541	543	545	547	549	551	553	555	557	559	561	563	565	567	569	571	573	575	577	579	581	583	585	587	589	591	593	595	597	599	601	603	605	607	609	611	613	615	617	619	621	623	625	627	629	631	633	635	637	639	641	643	645	647	649	651	653	655	657	659	661	663	665	667	669	671	673	675	677	679	681	683	685	687	689	691	693	695	697	699	701	703	705	707	709	711	713	715	717	719	721	723	725	727	729	731	733	735	737	739	741	743	745	747	749	751	753	755	757	759	761	763	765	767	769	771	773	775	777	779	781	783	785	787	789	791	793	795	797	799	801	803	805	807	809	811	813	815	817	819	821	823	825	827	829	831	833	835	837	839	841	843	845	847	849	851	853	855	857	859	861	863	865	867	869	871	873	875	877	879	881	883	885	887	889	891	893	895	897	899	901	903	905	907	909	911	913	915	917	919	921	923	925	927	929	931	933	935	937	939	941	943	945	947	949	951	953	955	957	959	961	963	965	967	969	971	973	975	977	979	981	983	985	987	989	991	993	995	997	999	1001	1003	1005	1007	1009	1011	1013	1015	1017	1019	1021	1023	1025	1027	1029	1031	1033	1035	1037	1039	1041	1043	1045	1047	1049	1051	1053	1055	1057	1059	1061	1063	1065	1067	1069	1071	1073	1075	1077	1079	1081	1083	1085	1087	1089	1091	1093	1095	1097	1099	1101	1103	1105	1107	1109	1111	1113	1115	1117	1119	1121	1123	1125	1127	1129	1131	1133	1135	1137	1139	1141	1143	1145	1147	1149	1151	1153	1155	1157	1159	1161	1163	1165	1167	1169	1171	1173	1175	1177	1179	1181	1183	1185	1187	1189	1191	1193	1195	1197	1199	1201	1203	1205	1207	1209	1211	1213	1215	1217	1219	1221	1223	1225	1227	1229	1231	1233	1235	1237	1239	1241	1243	1245	1247	1249	1251	1253	1255	1257	1259	1261	1263	1265	1267	1269	1271	1273	1275	1277	1279	1281	1283	1285	1287	1289	1291	1293	1295	1297	1299	1301	1303	1305	1307	1309	1311	1313	1315	1317	1319	1321	1323	1325	1327	1329	1331	1333	1335	1337	1339	1341	1343	1345	1347	1349	1351	1353	1355	1357	1359	1361	1363	1365	1367	1369	1371	1373	1375	1377	1379	1381	1383	1385	1387	1389	1391	1393	1395	1397	1399	1401	1403	1405	1407	1409	1411	1413	1415	1417	1419	1421	1423	1425	1427	1429	1431	1433	1435	1437	1439	1441	1443	1445	1447	1449	1451	1453	1455	1457	1459	1461	1463	1465	1467	1469	1471	1473	1475	1477	1479	1481	1483	1485	1487	1489	1491	1493	1495	1497	1499	1501	1503	1505	1507	1509	1511	1513	1515	1517	1519	1521	1523	1525	1527	1529	1531	1533	1535	1537	1539	1541	1543	1545	1547	1549	1551	1553	1555	1557	1559	1561	1563	1565	1567	1569	1571	1573	1575	1577	1579	1581	1583	1585	1587	1589	1591	1593	1595	1597	1599	1601	1603	1605	1607	1609	1611	1613	1615	1617	1619	1621	1623	1625	1627	1629	1631	1633	1635	1637	1639	1641	1643	1645	1647	1649	1651	1653	1655	1657	1659	1661	1663	1665	1667	1669	1671	1673	1675	1677	1679	1681	1683	1685	1687	1689	1691	1693	1695	1697	1699	1701	1703	1705	1707	1709	1711	1713	1715	1717	1719	1721	1723	1725	1727	1729	1731	1733	1735	1737	1739	1741	1743	1745	1747	1749	1751	1753	1755	1757	1759	1761	1763	1765	1767	1769	1771	1773	1775	1777	1779	1781	1783	1785	1787	1789	1791	1793	1795	1797	1799	1801	1803	1805	1807	1809	1811	1813	1815	1817	1819	1821	1823	1825	1827	1829	1831	1833	1835	1837	1839	1841	1843	1845	1847	1849	1851	1853	1855	1857	1859	1861	1863	1865	1867	1869	1871	1873	1875	1877	1879	1881	1883	1885	1887	1889	1891	1893	1895	1897	1899	1901	1903	1905	1907	1909	1911	1913	1915	1917	1919	1921	1923	1925	1927	1929	1931	1933	1935	1937	1939	1941	1943	1945	1947	1949	1951	1953	1955	1957	1959	1961	1963	1965	1967	1969	1971	1973	1975	1977	1979	1981	1983	1985	1987	1989	1991	1993	1995	1997	1999	2001	2003	2005	2007	2009	2011	2013	2015	2017	2019	2021	2023	2025	2027	2029	2031	2033	2035	2037	2039	2041	2043	2045	2047	2049	2051	2053	2055	2057	2059	2061	2063	2065	2067	2069	2071	2073	2075	2077	2079	2081	2083	2085	2087	2089	2091	2093	2095	2097	2099	2101	2103	2105	2107	2109	2111	2113	2115	2117	2119	2121	2123	2125	2127	2129	2131	2133	2135	2137	2139	2141	2143	2145	2147	2149	2151	2153	2155	2157	2159	2161	2163	2165	2167	2169	2171	2173	2175	2177	2179	2181	2183	2185	2187	2189	2191	2193	2195	2197	2199	2201	2203	2205	2207	2209	2211	2213	2215	2217	2219	2221	2223	2225	2227	2229	2231	2233	2235	2237	2239	2241	2243	2245	2247	2249	2251	2253	2255	2257	2259	2261	2263	2265	2267	2269	2271	2273	2275	2277	2279	2281	2283	2285	2287	2289	2291	2293	2295	2297	2299	2301	2303	2305	2307	2309	2311	2313	2315	2317	2319	2321	2323	2325	2327	2329	2331	2333	2335	2337	2339	2341	2343	2345	2347	2349	2351	2353	2355	2357	2359	2361	2363	2365	2367	2369	2371	2373	2375	2377	2379	2381	2383	2385	2387	2389	2391	2393	2395	2397	2399	2401	2403	2405	2407	2409	2411	2413	2415	2417	2419	2421	2423	2425	2427	2429	2431	2433	2435	2437	2439	2441	2443	2445	2447	2449	2451	2453	2455	2457	2459	2461	2463	2465	2467	2469	2471	2473	2475	2477	2479	2481	2483	2485	2487	2489	2491	2493	2495	2497	2499	2501	2503	2505	2507	2509	2511	2513	2515	2517	2519	2521	2523	2525	2527	2529	2531	2533	2535	2537	2539	2541	2543	2545	2547	2549	2551	2553	2555	2557	2559	2561	2563	2565	2567	2569	2571	2573	2575	2577	2579	2581	2583	2585	2587	2589	2591	2593	2595	2597	2599	2601	2603	2605	2607	2609	2611	2613	2615	2617	2619	2621	2623	2625	2627	2629	2631	2633	2635	2637	2639	2641	2643	2645	2647	2649	2651	2653	2655	2657	2659	2661	2663	2665	2667	2669	2671	2673	2675	2677	2679	2681	2683	2685	2687	2689	2691	2693	2695	2697	2699	2701	2703	2705	2707	2709	2711	2713	2715	2717	2719	2721	2723	2725	2727	2729	2731	2733	2735	2737	2739	2741	2743	2745	2747	2749	2751	2753	2755	2757	2759	2761	2763	2765	2767	2769	2771	2773	2775	2777	2779	2781	2783	2785	2787	2789	2791	2793	2795	2797	2799	2801	2803	2805	2807	2809	2811	2813	2815	2817	2819	2821	2823	2825	2827	2829	2831	2833	2835	283
------------	--	--	--	--	--	--	--	--	--	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----

[illegible]

[illegible]

[illegible]

[illegible]

GLOSSARY

5-HT = serotonin
 5-HT₁ = serotonin 1 receptor
 5-HT_{1A} = serotonin 1A receptor
 5-HT_{1B} = serotonin 1B receptor
 5-HT_{2A/2B} = serotonin 2A and serotonin 2B receptor
 5-HT_{2B} = serotonin 2B receptor
 5-HT_{2C} = serotonin 2C receptor
 5-HT₃ = serotonin 3 receptor
 5-HT₄ = serotonin 4 receptor
 AMPA = alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionate
 CB1 = cannabinoid receptor 1
 COX-2 = COX-inhibiting nitric oxide donator
 COX = cyclooxygenase
 COX-2 = cyclooxygenase 2
 CRF-1 = Corticotropin-Releasing Factor Receptor 1
 D1 = Dopamine 1
 D2 = Dopamine 2
 D3 = Dopamine 3
 D4 = Dopamine
 ERK = extracellular signal-related kinase
 GABA = gamma-aminobutyric acid
 GABA-A = gamma-aminobutyric acid A receptor
 GABA-B = gamma-aminobutyric acid B receptor
 GPCR = G-Protein-Coupled Receptor
 GR = glucocorticoid receptor
 H3 = histamine H3-receptor
 MAO = mono-amine oxidase
 MAO-A = mono-amine oxidase A
 MAO-B = mono-amine oxidase B
 MCA = melanocortin-4 receptor
 MCH = Melanin concentrating hormone
 mGluR = metabotropic glutamate receptor
 mGluR1 = metabotropic receptor
 NAR1 = selective nor-adrenaline re-uptake inhibitor
 nNOS = nitroendophectic neurotoxic antidepressant
 NDRI = selective nor-adrenaline and dopamine re-uptake inhibitor
 NGF = Nerve Growth Factor
 NGF = nerve growth factor
 NGF = neurotrophin 1 receptor
 NGF = neurotrophin 2 receptor
 NGF = neurotrophin 3 receptor
 NMDA = N-Methyl-D-aspartate
 NSAID = Non-steroidal anti-inflammatory drugs
 PDE4 = phosphodiesterase-4
 RIMA = reversible inhibitor of mono-amine oxidase A
 SCD-11 = G protein-coupled receptor
 SDA = Serotonin/Dopamine Antagonist
 SDRI = selective serotonin and dopamine reuptake inhibitor
 SMDRI = selective serotonin, nor-adrenaline and dopamine reuptake inhibitor
 SNRI = selective serotonin and nor-adrenaline reuptake inhibitor
 SSRI = selective serotonin reuptake inhibitor
 VIB = vasopressin 1B receptor

PHARMAC GROUP (see overview hereunder)	IR, PH, PROF.	PHARMACOLOGICAL PROFILE	MAIN INDICATIONS	COMPOUND	DOSE RANGE	COMPANY
Monoaminergic Transmitter Systems	1	5-HT reuptake enhancer	Depression / Anxiety	THW-309	25 to 50mg daily	Phar
	2	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	Phar
	3	5-HT _{1A} agonist	Depression / Anxiety	THW-309	150mg to 450mg	MediNova
	4	5-HT _{1A} agonist	Depression / Anxiety	THW-309	20mg to 50mg	MediNova
	5	5-HT _{1A} agonist	Depression / Anxiety	THW-309	1 to 2mg daily	MediNova
	6	5-HT _{1A} agonist	Depression / Anxiety	THW-309	unknown	MediNova
	7	5-HT _{1A} agonist	Depression / Anxiety	THW-309	unknown	MediNova
	8	5-HT _{1A} agonist	Depression / Anxiety	THW-309	unknown	MediNova
	9	5-HT _{1A} agonist	Depression / Anxiety	THW-309	unknown	MediNova
	10	5-HT _{1A} agonist	Depression / Anxiety	THW-309	unknown	MediNova
Monoaminergic Transmitter Systems	11	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	12	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	13	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	14	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	15	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	16	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	17	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	18	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	19	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	20	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
Monoaminergic Transmitter Systems	21	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	22	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	23	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	24	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	25	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	26	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	27	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	28	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	29	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	30	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
Monoaminergic Transmitter Systems	31	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	32	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	33	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	34	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	35	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	36	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	37	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	38	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	39	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	40	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
Monoaminergic Transmitter Systems	41	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	42	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	43	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	44	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	45	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	46	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	47	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	48	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	49	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	50	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
Monoaminergic Transmitter Systems	51	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	52	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	53	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	54	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova
	55	5-HT _{1A} antagonist	Depression / Anxiety	THW-309	unknown	MediNova

PHARMAC. GROUP (see overview hereunder)	nr. PH. PROF.	PHARMACOLOGICAL PROFILE	MAIN INDICATIONS	COMPOUND	DOSE RANGE	COMPANY
Pedagogic Mechanisms of Domestics of the Alzheimer Type	16 amyloidase 17 amyloid aggregation-inhibitor		ADHD	methylphenidate transdermal system		Novon Pharmaceuticals
Endocrine System	18 amyloid aggregation-inhibitor		Alzheimer's Disease	Ashford	200 to 30mg daily	Harscochem
Monoaminergic Transmitter Systems	19 amyloid aggregation-modulator		Alzheimer's Disease	JAN		Pharmacia
Other Unknown	19 beta 3 adrenoceptor agonist		Female Sexual Dysfunction	LX02226		Liquid Pharmaceuticals
	20 GABA Channel Modulator		Depression / Anxiety	SR 58811	unknown	Sandoz Synthabo
	21 glutamate receptor antagonist		Alzheimer's Disease	MEM 1003		Novon Pharmaceuticals
	22 calcium K ⁺ inhibitor		Parkinson's Disease	leflunomide	unknown	Novon Pharmaceuticals
	23 alpha uptake enhancer		Schizophrenia	SR 141716		Sandoz Synthabo
	24 COX-2 inhibitor		Pain	AG2786		Ciba-Sandoz
	COX-2 inhibitor		Alzheimer's Disease	MUC-21	20 to 160mg daily	Merck Serenel Pharmas
	COX-2 inhibitor		Pain	rofecoxib		Pfizer
	COX-2 inhibitor		Pain	valdecoxib		Pfizer
	COX-2 inhibitor		Pain	etoricoxib		Merck
	COX-2 inhibitor		Pain	COX 188	20 to 120mg daily	Novartis Pharmaceuticals
	COX-2 inhibitor		Pain	peroxob	100 to 800mg daily	Pfizer
	COX-2 inhibitor		Pain	ART-663	20 to 100mg daily	Abbott
	COX-2 inhibitor		Pain	AZD3582		AstraZeneca
	COX-2 inhibitor		Pain	ZD7417	375mg daily	Novartis Pharmaceuticals
	25 COX-inhibiting anti-oxidant donors (CHODs) COX-inhibiting nitric oxide donors (CINODs)		Depression	MAE 861	unknown	Rice
	GRF1 antagonist		Depression / Anxiety	R121819	5 to 80mg daily	Novartis Pharmaceuticals
	GRF1 antagonist		Depression / Anxiety	dexameth	unknown	Pfizer
	GRF1 antagonist		Depression	TZSC20		GlenSmithKline
	GRF1 antagonist		Depression / Anxiety	NE-39041		Neurocrine Biosciences
	GRF1 antagonist		Depression / Anxiety	CP-142428		Pfizer
	GRF1 antagonist		Depression / Anxiety	CP-448,187		Dag Abnata Sciences
	GRF1 antagonist		Cocaine Dependence	DAS-431	unknown	Off robot
	27 O1 receptor agonist		Schizophrenia	amiphenide		Sohay
	28 O2 receptor antagonist		Schizophrenia	afeganor		? ?
	O2 receptor antagonist		Schizophrenia	BSE-201840		? ?
	O2 antagonist		Cocaine Dependence	PD 58491		? ?
	O2 antagonist		Cocaine Dependence	BSE-201840		? ?
	O2 antagonist		Parkinson's Disease	PD 58491		? ?
	O2 antagonist		Parkinson's Disease	BSE-201840		? ?
	O2 antagonist		Schizophrenia	PD 58491		? ?
	O2 antagonist		Cocaine Dependence	GBR 12869		National Institute on Drug Abuse
	30 DA uptake reboar		Parkinson's Disease	fenamitab		Novon Pharmaceuticals
	DA uptake reboar		Parkinson's Disease	emamitab		Pfizer
	31 fosapamine agonist		Parkinson's Disease	mepazine CDS (Once-a-Day Transdermal Patch)	4 to 16mg daily	Schering Pharma
	fopamine agonist		Parkinson's Disease	nopolone HCL (contraindicated formulation)	4.5 to 13.5mg daily	GlenSmithKline
	fopamine agonist		Cocaine Dependence	cabaprine	0.75 to 24mg daily	Abbott
	fopamine agonist		Cocaine Dependence	carbamaz		ELUO Pharmaceuticals
	fopamine agonist		Parkinson's Disease	contamab		Lepra
	fopamine agonist		Parkinson's Disease	DOB-57		Wegh
	fopamine agonist		Parkinson's Disease, Comorbid	SLV 308		Sohay
	fopamine agonist		Depression / Anxiety	SD2504		Santal
	fopamine agonist		Parkinson's Disease	SD2504		Service
	fopamine agonist		Parkinson's Disease	homocysteine		Novartis Pharmaceuticals
	fopamine agonist		Parkinson's Disease	choleste		ALM - Not From Backer (?)

PHARMAC. GROUP (see overview hereunder)	RT. PH. PROF.	PHARMACOLOGICAL PROFILE	MAIN INDICATIONS	COMPOUND	DOSE RANGE	COMPANY
Enzyme System	32	GRK activation	Alzheimer's Disease	CP-115,893	50 to 100mg daily	Center Pharmaceuticals
Inhibitory Amino Acid System	33	GABA-A agonist	Alzheimer's Disease	Nefazodone	unknown	Daini Seiyaku, JPN/Boehringer, BRD
Inhibitory Amino Acid System	34	GABA-A agonist	Alzheimer's Disease	Gabaudin	5 to 20mg daily	Lockwood
Inhibitory Amino Acid System	35	GABA-A agonist	Alzheimer's Disease	extastodine	2 to 3mg daily	Sepacor
Inhibitory Amino Acid System	36	GABA-A antagonist	Alzheimer's Disease	Zapfen MR sustained-release version	10 to 20mg daily	Sandoz-Synthelabo
Inhibitory Amino Acid System	37	GABA-A antagonist	Alzheimer's Disease	indolone	10 to 20mg daily	DOV/Neurocrine
Inhibitory Amino Acid System	38	GABA-A antagonist	Alzheimer's Disease	pagodone	30mg daily	Indevus
Inhibitory Amino Acid System	39	GABA-A antagonist	Alzheimer's Disease	Zapfen extended-release	10mg daily	King Pharmaceuticals
Inhibitory Amino Acid System	40	GABA-A antagonist	Alzheimer's Disease	SEP174559	10mg daily	Sepacor
Inhibitory Amino Acid System	41	GABA-A antagonist	Alzheimer's Disease	SL 65,1430	10 to 60mg daily	Sandoz-Synthelabo
Inhibitory Amino Acid System	42	GABA-A antagonist	Alzheimer's Disease	CP-720,330 (NCD 95-3)	10 to 60mg daily	Neurocrine
Inhibitory Amino Acid System	43	GABA-A antagonist	Alzheimer's Disease	NCD 95-3	10 to 60mg daily	DOV
Inhibitory Amino Acid System	44	GABA-A antagonist	Alzheimer's Disease	AVE 7398	15mg daily	Aventis
Inhibitory Amino Acid System	45	GABA-A antagonist	Alzheimer's Disease	GNF	15mg daily	National Institute on Drug Abuse
Inhibitory Amino Acid System	46	GABA-A antagonist	Alzheimer's Disease	LY35140	15mg daily	El Lilly
Inhibitory Amino Acid System	47	GABA-A antagonist	Alzheimer's Disease	RY204	500 to 1200mg daily	Roche
Inhibitory Amino Acid System	48	GABA-A antagonist	Alzheimer's Disease	Mefenitone	500 to 1200mg daily	Concept
Inhibitory Amino Acid System	49	GABA-A antagonist	Alzheimer's Disease	ORG M1704659	unknown	Organon
Inhibitory Amino Acid System	50	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	51	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	52	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	53	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	54	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	55	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	56	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	57	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	58	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	59	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	60	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	61	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	62	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	63	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	64	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	65	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	66	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	67	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	68	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	69	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	70	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	71	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	72	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	73	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	74	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	75	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	76	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	77	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	78	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	79	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	80	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	81	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	82	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	83	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	84	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	85	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	86	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	87	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	88	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	89	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	90	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	91	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	92	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	93	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	94	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	95	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	96	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	97	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	98	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	99	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott
Inhibitory Amino Acid System	100	GABA-A antagonist	Alzheimer's Disease	ABT 534	unknown	Abbott

[illegible]

PHARMAC. GROUP (see overview hereunder)	nr. PH. PROF.	PHARMACOLOGICAL PROFILE	MAIN INDICATIONS	COMPOUND	DOSE RANGE	COMPANY
Monoaminergic Transmitter Systems	61	5-HT _{1C} modulation	Depression	SNR-2		Synapse
	64	5-HT _{1A} SDA	Schizophrenia	SNR-2		AstraZeneca
		SDA	Schizophrenia	SNR-2		Bristol-Myers Squibb
		SDA	Schizophrenia	SNR-2		Johnson & Johnson Pharmaceuticals
		SDA	Schizophrenia	SNR-2		Kiwi / BASF
		SDA	Schizophrenia	SNR-2		Lilly
		SDA	Schizophrenia	SNR-2		Novartis Pharmaceuticals
		SDA	Schizophrenia	SNR-2		Pfizer
		SDA	Schizophrenia	SNR-2		Roche
		SDA	Schizophrenia	SNR-2		Schering
Monoaminergic Transmitter Systems	62	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	63	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	64	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	65	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	66	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	67	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	68	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	69	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	70	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	71	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
Monoaminergic Transmitter Systems	72	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	73	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	74	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	75	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	76	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	77	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	78	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	79	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	80	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	81	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
Monoaminergic Transmitter Systems	82	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	83	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	84	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	85	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	86	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	87	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	88	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	89	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	90	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	91	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
Monoaminergic Transmitter Systems	92	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	93	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	94	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	95	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	96	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	97	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	98	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	99	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	100	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering
	101	5-HT _{1A} SDA	Schizophrenia	SNR-2		Schering

PHARMAC. GROUP (see overview hereunder)	nr. PH. PROF.	PHARMACOLOGICAL PROFILE	MAIN INDICATIONS	COMPOUND	DOSE RANGE	COMPANY
Neurotransmitter Transmitter Systems	57	SSRI	Depression / Anxiety	paroxetine	10 to 20mg daily	GlaxoSmithKline
	58	SSRI	Depression / Anxiety	citalopram	10 to 40 mg daily	Lundbeck / Forest Laboratories
	59	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	of patient
	60	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	of patient
	61	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Pfizer
	62	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Sandoz-Synthelabo
	63	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Ferraro
	64	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	65	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	66	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	67	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	68	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
Peptide Transmitter Systems	69	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	70	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	71	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	72	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	73	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	74	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	75	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	76	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	77	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	78	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	79	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	80	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
Other / Unknown	81	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	82	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	83	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	84	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	85	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	86	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	87	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	88	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	89	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	90	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	91	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	92	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
Other / Unknown	93	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	94	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	95	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	96	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	97	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	98	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	99	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	100	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	101	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	102	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	103	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals
	104	SSRI	Depression / Anxiety	fluoxetine	10 to 40 mg daily	Novartis Pharmaceuticals

Examples

Example 1: Measuring pKi values of test compounds

5 In Table 1, the pKi values of test compounds are given for each of the dopamine receptors, 5HT receptors, adrenergic receptors and the histamine1 receptor. The affinity of test compounds for the respective receptors has been performed according to conventional procedures known in the art.

10 An indication "0" means that no affinity has been measured between the test compound and the receptor.

The columns displaying the pKi values for the D4 and the 5-HT2A receptor are filled with dark grey. pKi values between 8 and 9 and higher than 9 are represented by light grey boxes.

15 Example 2: Foregoing pipamperon-citalopram treatment in major depressive disorder: a placebo and active controlled period finding clinical trial

Table 2 represents the set-up of a clinical trial comprising for treatment groups:

20 Group Plc – Active / Day 0 represents the group receiving 10 mg citalopram, twice a day, starting the first day (Day 0) of active treatment in the clinical trial. This administration regime is also indicated as the mono therapy.

Group Pip - Active / Day 0 represents the group receiving a combination of 4 mg pipamperon and 10 mg citalopram, twice a day, starting the first day (Day 0) of active treatment in the clinical trial. This administration regime is also indicated as the non-foregoing combo therapy.

25 Group Pip - Active / Day 4 represents the group receiving 4 mg pipamperon, twice a day, starting the first day (Day 0) of active treatment in the clinical trial, followed by a combination of 4 mg pipamperon and 10 mg citalopram, twice a day, starting the fifth (Day 4) day of active treatment in the clinical trial. This administration regime is also indicated as the foregoing therapy with combination therapy starting after 4 days of active treatment.

30 Group Pip - Active / Day 7 represents the group receiving 4 mg pipamperon, twice a day, starting the first day (Day 0) of active treatment in the clinical trial, followed by a combination of 4 mg pipamperon and 10 mg citalopram, twice a day, starting the eight (Day 7) day of

active treatment in the clinical trial. This administration regime is also indicated as the foregoing therapy with combination therapy starting after 7 days of active treatment.

All subjects also undergo a placebo (PLC) run-in therapy, administered during a period of about 7 days before the active treatment starts.

5 During daily (D), weekly (W) or monthly (M) visits, several parameters are measured.

Under NECT is to be understood: Neuronal E-clinical Trial = Vesalius Expert development for this trial which includes the bottom-up measurement of:

- In- and exclusion-criteria
- Functional status evaluation
- 10 - Medical history
- (Pre-)treatment signs & symptoms
- DSM-IV rules for diagnosis & efficacy
- HDRS-28 (Hamilton Depression Rating Scale - 28 items)
- Medical resource utilisation
- 15 - Pre-trial & Concomittant medication
- Drug administration
- (Serious) Adverse events
- Admission to the acute and extension phase of treatment
- Right flow of the trial
- 20

Example 3: combo pipamperon-citalopram: therapeutic use in Major Depression

25 Purpose

Pipamperon (1'-[3-(p-Fluorobenzoyl)propyl]-[1,4'-bipiperidine]-4'-carboxamide), the active ingredient of Dipiperon (Janssen-Cilag B.V), administered to patients in a dose ranging between 8 and 12 mg is claimed via its specific pharmacological properties to be a booster of the antidepressant effect of the selective serotonin re-uptake inhibitor

30 citalopram. Preferably, pipamperon is administered daily at least 4-5 days before administering said antidepressant. The mechanism of boosting of pipamperon has to deal with (i) the selective affinity for the dopamine-4 (D4) receptor with a pKi value equal to or higher than 8 towards the D4 receptor and less than 8 towards other dopamine receptors, and (ii) the selective affinity for the 5-HT2A receptor with a pKi value equal to or higher

35 than 8 towards the 5-HT2A receptor and less than 8 towards other 5HT receptors. This semi-naturalistic open label study investigated the efficacy and tolerability of the combo pipamperon - citalopram in the treatment of patients with major depression.

Details

Design: Semi-naturalistic i.e. inclusion of every 'natural' patient in an outpatient practice but without concomitant use of mood enhancing drugs, open label

Control: No

Phase: Phase IIa – preliminary Proof of Concept

Location: Belgium – Research Centre ANIMA, Alken

End Points : Assessment scale scores, Hamilton Depression Rating Scale 17 items, Reduction, Response, Remission

Medication: Exclusion of mood stabilisers, antipsychotics (typical and atypical) and other antidepressants

Subjects

Type	No.	Sex	Age
Patients	23	10 male & 13 female	23-80 (mean 47) years

- 5 Characteristics: patients had a major depressive disorder according to DSM-IV criteria, with or without a chronic course and a treatment refractory state towards another SSRI then citalopram.

Treatments

- 10 PIP-CIT¹ add-on: citalopram from day minus 60-20 – pipamperon from DAY 0

Drug/Treatment	Dose	Route	Frequency	Duration
Pipamperon ¹	+ Pip.: 8-12 mg/day – Cit.: 20-	PO	bid	8 weeks
Citalopram ¹	40 mg/day			

1. Pipamperon (Pip) and citalopram (Cit) dosage was adjusted according to clinical response.

15

PIP-CIT¹ fore-going 1-5: pipamperon from day 0 – cital from day 1-5

Drug/Treatment	Dose	Route	Frequency	Duration
Pipamperon ¹	+ Pip.: 8-12 mg/day – Cit.: 20-	PO	bid	8 weeks
Citalopram ¹	40 mg/day			

- 20 1. Pipamperon (Pip) and citalopram (Cit) dosage was adjusted according to clinical response.

25

PIP-CIT¹ fore-going 6-8: pipamperon from day 0 – citalopram from day 6-8

193

Drug/Treatment	Dose	Route	Frequency	Duration
Pipamperon ¹	+ Pip.: 8-12 mg/day – Cit.: 20-	PO	bid	8 weeks
Citalopram ¹	40 mg/day			

1. Pipamperon (Pip) and citalopram (Cit) dosage was adjusted according to clinical response.

5 Results

Mean Medication	Used	PIP-CIT add-on	PIP-CIT foregoing
		After 20-60 DAYS (mean 33) (n = 5)	1-5 DAYS (mean 4) (n = 15) 6-8 DAYS (mean 7) (n = 3)
Pipamperone		9mg/day	10mg/day
Citalopram		30mg/day	26mg/day
Depression scale scores			
HDRS 17-item total score			
baseline		29	23
endpoint (week 8)		4	5
diminishment at week 8		-25 (+8/-9)	-18 (+8/-8)
% reduction at week 8		86 (+14/-12)	80 (+20/-30)
response ¹ at week 8		5 (100%)	15 (100%)
remission ² at week 8		4 (80%)	10 (67%)

1. Response = $\geq 50\%$ reduction in HDRS 17-item score;

2. Remission = HDRS 17-item score < 8

- 10 Notably, the results obtained are highly significant since the variability in every group is distributed evenly around the mean.

Add-on PIP-CIT

- 15 Figure 1 schematically depicts the "add-on" treatment with pipamperon 8-12 (mean 9) mg (bid) after treatment with citalopram 10-20 (mean 30) mg (bid) during 20-60 (mean 33) days (PIPCIT ADD-ON) with HDRS-17. Totalscore is 29 at baseline in MDD in comparison with the standard efficacy of antidepressants in clinical trials according to Khan *et al.* (2000), in "Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical Trials" (Arch. of General Psychiatry, Vol. 57, April 2000).

Figure 2 schematically depicts the HDRS-17 change from baseline in the combo pipamperon as "add-on" to citalopram vs SNRI (duloxetine) in Major Depression. Treatment with pipamperon 8-12 (mean 9 mg/day) during 20-60 (mean 33) days after treatment with SSRI (n=5). The SNRI (duloxetine) treatment was 40-120 mg/day (n = 152) according to Goldstein *et al.*, (Clin. Psychiatry, in press).

Figure 3 schematically depicts the remission rates (HDRS-17 ≤ 7) with the combo pipamperon as "add-on" to citalopram vs SNRI (venlafaxine) vs SSRIs vs placebo in Major Depression. Treatment with pipamperon 8-12 (mean 9 mg/day) during 20-60 (mean 33) days after treatment with SSRI (n=5). Treatment with the SNRI venlafaxine is according to a meta-analysis of Thase *et al.* (Br. J. Psychiatry (2001) 178:234-241). Treatment with SSRIs is according to a meta-analysis of Thase *et al.* (Br. J. Psychiatry (2001) 178:234-241). Treatment with placebo is according to a meta-analysis of Thase *et al.* (Br. J. Psychiatry (2001) 178:234-241).

Fore-going 1-5 PIP-CIT

Figure 4 schematically depicts the "fore-going" treatment during 1-5 (mean 4) days with pipamperon 8-12 (mean 10) mg (bid), followed with the combination treatment of pipamperon and citalopram 20-50 (mean 26) mg/day (bid) (PIPCIT FG 1-5) in MDD (HDRS-17 at BL = 23) in comparison with the standard efficacy of antidepressants in clinical trials according to Khan *et al.* (2000), in "Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical Trials" (Arch. of General Psychiatry, Vol. 57, April 2000).

Figure 5 schematically depicts the HDRS-17 change from baseline in the combo pipamperon-citalopram treatment with a "fore-going" treatment of 4 days with pipamperon (10 mg/day) vs SNRI (duloxetine) in Major Depression. Treatment with the combo pipamperon-citalopram with pipamperon 8-12 (mean 10 mg/day) (bid) 1-5 (mean 4) days before treatment with SSRI (n=15). The SNRI (duloxetine) treatment was 40-120 mg/day (n = 152) according to Goldstein *et al.*, (Clin. Psychiatry, in press).

Figure 6 schematically depicts the remission rates (HDRS-17 ≤ 7) with the combo pipamperon with a "fore-going" treatment of 4 days with pipamperon (10 mg/day) vs SNRI (venlafaxine) in Major Depression. Treatment with the combo pipamperon-citalopram was with pipamperon 8-12 (mean 10 mg/day) during 1-5 (mean 4) days before treatment with the

SSRI (n=5). Treatment with the SNRI venlafaxine is according to a meta-analysis of Thase et al. (Br. J. Psychiatry (2001) 178:234-241). Treatment with SSRIs is according to a meta-analysis of Thase et al. (Br. J. Psychiatry (2001) 178:234-241). Treatment with placebo is according to a meta-analysis of Thase et al. (Br. J. Psychiatry (2001) 178:234-241).

5

Fore-going 6-8 PIP-CIT

Figure 7 schematically depicts the "fore-going" treatment during 6-8 (mean 7) days with pipamperon 8-12 (mean 11) mg/day (bid), followed with the combination treatment of pipamperon and citalopram 20-40 (mean 30) mg/day (bid) (PIPCIT FG 6-8) in MDD (HRS-17 at BL = 28) in comparison with the standard efficacy of antidepressants in clinical trials according to Khan et al. (2000), in "Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical Trials" (Arch. of General Psychiatry, Vol. 57, April 2000).

Figure 8 schematically depicts the HRS-17 change from baseline in the combo pipamperon-citalopram treatment with a "fore-going" treatment of 7 days with pipamperon (11 mg/day) vs SNRI (duloxetine) in Major Depression. Treatment with the combo pipamperon-citalopram with pipamperon 8-12 (mean 11 mg/day) (bid) 6-8 (mean 7) days before treatment with SSRI (n=3). The SNRI (duloxetine) treatment was 40-120 mg/day (n = 152) according to Goldstein et al., (Clin. Psychiatry, in press).

Comparison "add-on" vs "fore-going"

Figure 9 schematically depicts a comparison between "fore-going" and "add-on" treatments with pipamperon (8-12 mg/day; bid) and citalopram (20-40 mg/day; bid) in MDD in comparison with the standard efficacy of antidepressants in clinical trials according to Khan et al. (2000), in "Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical Trials" (Arch. of General Psychiatry, Vol. 57, April 2000).

Figure 10 schematically depicts a comparison between "fore-going" and "add-on" treatments. In particular, the HRS-17 change from baseline between "fore-going" and "add-on" treatment with pipamperon (8-12 mg/day; bid) and citalopram (20-40 mg/day; bid) in comparison with the SNRI duloxetine in Major Depression is depicted. Treatment with the combo pipamperon as "add-on" to citalopram, with pipamperon 8-12 mg/day (mean 9 mg/day) 20-60 (mean 33) days after treatment with the SSRI (n=5). Treatment with the combo pipamperon-citalopram, with pipamperon 8-12 mg/day (mean 11 mg/day; bid) 6-8 days (mean 7 days) before treatment with the SSRI (n = 15). Treatment with the combo

pipamperon-citalopram, with pipamperon 8-12 mg/day (mean 10 mg/day; bid) 1-5 days (mean 4 days) before treatment with the SSRI (n = 15). The SNRI (duloxetine) treatment was 40-120 mg/day (n = 152) according to Goldstein *et al.*, (Clin. Psychiatry, In press).

- 5 Figure 11 schematically depicts the remission rates (HDRS-17 ≤ 7) in a comparison between "fore-going" and "add-on" treatment with pipamperon (8-12 mg/day; bid) and citalopram (20-40 mg/day; bid) in comparison with the SNRI venlafaxine in Major Depression. Treatment with the combo pipamperon-citalopram was with pipamperon 8-12 (mean 10 mg/day) during 1-5 (mean 4) days before treatment with the SSRI (n = 15).
- 10 Treatment with the SNRI venlafaxine is according to a meta-analysis of Thase *et al.* (Br. J. Psychiatry (2001) 178:234-241). Treatment with pipamperon as "add-on" to citalopram, with pipamperon 8-12 (mean 9 mg/day) during 20-60 (mean 33) days after treatment with SSRI (n = 5).
- 15 The intention-to-treat / last-observation-carried-forward analysis showed a high therapeutic efficacy according HDRS 17-item in all the treatment groups. This was especially true for the 'add-on' group probably caused by the longer treatment with an active antidepressant (+33 days). The huge therapeutic effect observed in the 'PIP-CIT 1-5' group present for at a mean dosage of pipamperon of 10 mg per day and administered
- 20 the first four days of treatment without an active antidepressant, indicates the boosting effect of pipamperon on the SSRI citalopram at an extremely and thus unconventional low dose. Only 1 patient discontinued treatment due to a lost of follow-up.

Adverse Events

Side effects (patients)	PIP-CIT	add-on	PIP-CIT foregoing
	After 20-60 DAYS (mean 33) (n = 5)	1-5 DAYS (mean 4) (n = 15)	6-8 DAYS (mean 7) (n = 3)
Discontinued treatment due to adverse events	0	0	0
By system:			
body as a whole	0	0	0
central and peripheral nervous system	1(20%)	4(26.6%)	0
gastrointestinal	1(20%)	5(33%)	2(66.6%)
musculoskeletal	1(20%)	3(20%)	0
psychiatric	0	0	0
respiratory	0	1(6.6%)	0
skin and appendages	1(20%)	2(13.3%)	1(33.3%)
vascular	0	1(6.6%)	0
urinary	0	1(6.6%)	0

Laboratory parameters, ECG, bodyweight and vital signs were not measured since this was a naturalistic study.

5

AssessmentOutcome

Efficacy: the 4-day fore-going combo pipamperon 8-12mg/d - citalopram 20-40mg/day is comparable to the add-on combo pipamperon-citalopram.

10 **Efficacy:** the 4-day fore-going combo pipamperon 8-12mg/d - citalopram 20-40mg/day is larger than the 7-day fore-going combo pipamperon 8-12mg/d - citalopram 20-40mg/day.

Efficacy: the combo pipamperon 8-12mg/d - citalopram 20-40mg/day is larger than the in the art known antidepressants SSRIs.

15 Tolerability

Tolerability: the 4-day fore-going treatment is comparable to the 7-day fore-going combo is comparable to add-on combo pipamperon-citalopram.

Tolerability: no discontinued treatment due to adverse events.

Study Messages

5 The boosting effect of pipamperon at an extremely unconventional low dose on a SSRI is indicated since the efficacy of the 'add-on' and '4-day fore-going' combo 'pipamperon 8 - 12 mg/d - citalopram 20 - 40 mg/day' is in this study as twice higher as known in the art in the treatment of patients with major depression.

10 The combo pipamperon-citalopram is generally well tolerated in patients with depression i.e. at least no specific added adverse events were occurring by adding pipamperon at the doses used in the study.

Example 4: combo pipamperon-citalopram: therapeutic use in Obsessive-Compulsive Disorder (OCD).

Purpose

5 Pipamperon (1'-[3-(p-Fluorobenzoyl)propyl]-[1,4'-bipiperidine]-4'-carboxamide), the active ingredient of Dipiperon (Janssen-Cilag B.V), administered to a patient in a dose ranging between 8 and 12 mg is claimed via its specific pharmacological properties to be a booster of the effect of the selective serotonin re-uptake inhibitor citalopram towards OCD. Preferably, pipamperon is administered daily at least 4-5 days before administering
10 said antidepressant. The mechanism of boosting of pipamperon has to deal with (i) the selective affinity for the dopamine-4 (D4) receptor with a pKi value equal to or higher than 8 towards the D4 receptor and less than 8 towards other Dopamine receptors, and (ii) the selective affinity for the 5-HT2A receptor with a pKi value equal to or higher than 8 towards the 5-HT2A receptor and less than 8 towards other 5HT receptors. This semi-
15 naturalistic open label study investigated the efficacy and tolerability of the combo pipamperon - citalopram in the treatment of patients with OCD.

Details

Design: Semi-naturalistic i.e. inclusion of every 'natural' patient in an outpatient practice but without concomitant use of mood enhancing drugs, open label

Control: No

Phase: Phase IIa – preliminary Proof of Concept

Location: Belgium – Research Centre ANIMA, Alken

End Points : Assessment scale scores, Yale-Brown Obsessive-Compulsive Scale, Reduction, Remission

Medication: Exclusion of mood stabilisers, antipsychotics (typical and atypical) and other antidepressants

20 **Subjects**

Type	No.	Sex	Age
Patients	7	1 male & 7 female	20-63 (mean 33) years

Characteristics: patients had an obsessive-compulsive disorder according to DSM-IV criteria, with or without a chronic course and a treatment refractory state towards another SSRI then citalopram.

200

TreatmentsPIP-CIT¹ ADD-ON: citalopram from DAY minus 730-60 – pipamperon from DAY 0

Drug/Treatment	Dose	Route	Frequency	Duration
Pipamperone ¹	+ Pip.: 8-16 mg/day – Cit.: 30-80	PO	bid	12 weeks
Citalopram ¹	mg/day			

- 5 1. Pipamperone (Pip) and Citalopram (Cit) dosage was adjusted according to clinical response.

PIP-CIT¹ FORE-GOING 4-6: pipamperon from DAY 0 – citalopram from DAY 4-6

10

Drug/Treatment	Dose	Route	Frequency	Duration
Pipamperone ¹	+ Pip.: 8-16 mg/day – Cit.: 30-80	PO	bid	12 weeks
Citalopram ¹	mg/day			

1. Pipamperone (Pip) and Citalopram (Cit) dosage was adjusted according to clinical response.

Results

		PIP-CIT add-on
		after 730-60 DAYS (mean 241) (n = 6)
		with mean Cit. 54mg/d and Pip. 11mg/d
		PIP-CIT foregoing
		4-6 DAYS (mean 5) (n = 2)
		with mean Cit. 60mg/d and Pip. 10mg/d
Y-BOCS score		
<u>Baseline</u>		
Total	31	
Obsessions	18	
Compulsions	13	
Endpoint (week 12)		
<u>Total</u>	<u>15</u>	
diminishment	-16 (+16/-11)	
% reduction	53	
<u>Obsessions</u>		
total	8	
diminishment	-10 (+9/-7)	
% reduction	57	
<u>Compulsions</u>		
total	7	
diminishment	-6 (+7/-6)	
% reduction	45	
<u>% Remission</u>		
YBOCS score ≤8	29	
BOCS score ≤16	57	

Notably, the results obtained are highly significant since the variability in every group is distributed evenly around the mean.

Figure 12 schematically depicts the Y-BOCS total score: "fore-going" and "add-on" treatment with pipamperon (8-15 mg/day; bid) and citalopram (30-80 mg/day; bid) in comparison with the SSRI fluvoxamine in OCD. Treatment with the combo pipamperon-citalopram (n = 7). Treatment with fluvoxamine (controlled release) mean 271 mg/day (n = 253) is according to Hollander et al. (2003).

Figure 13 schematically depicts the Y-BOCS obsession score: "fore-going" and "add-on" treatment with pipamperon (8-15 mg/day; bid) and citalopram (30-80 mg/day; bid) in comparison with the SSRI fluvoxamine in OCD. Treatment with the combo pipamperon-citalopram (n = 7). Treatment with fluvoxamine (controlled release) mean 271 mg/day (n = 253) is according to Hollander et al. (2003).

Figure 14 schematically depicts the Y-BOCS compulsion score: "fore-going" and "add-on" treatment with pipamperon (8-16 mg/day; bid) and citalopram (30-80 mg/day; bid) in comparison with the SSRI fluvoxamine in OCD. Treatment with the combo pipamperon-citalopram (n = 7). Treatment with fluvoxamine (controlled release) mean 271 mg/day (n = 253) is according to Hollander et al. (2003).

The intention-to-treat / last-observation-carried-forward analysis showed a high therapeutic efficacy according Y-BOCS total score, obsession and compulsion scores. This indicates the boosting effect of pipamperon on the SSRI citalopram at an extremely and thus unconventional low dose. No patient discontinued treatment.

Assessment

Efficacy: the combo pipamperone 8-16mg/d - citalopram 30-80mg/day > the in the art known compounds effective towards OCD (Hollander E, Koran LM, Goodman WK, Greist JH, Ninan PT, et al. A double-blind, placebo-controlled study of the efficacy and safety of controlled-release fluvoxamine in patients with obsessive-compulsive disorder. Journal of Clinical Psychiatry 64: 640-647, Jun 2003 Mount Sinai School of Medicine, New York, New York, USA; Solvay Pharmaceuticals Inc., Marietta, Georgia, USA).

Study Messages

The boosting effect of pipamperon at an extremely unconventional low dose on a SSRI is indicated since the efficacy of the 'add-on' and 'fore-going' combo 'pipamperon 8-15 mg/d - citalopram 30-80 mg/day' is in this study as twice higher as known in the art in the treatment of patients with obsessive-compulsive disorder.

Example 5: combo pipamperon-citalopram: therapeutic use in Panic Disorder.**Purpose**

Preliminary examination of a "fore-going" and "add-on" treatment with pipamperon and citalopram in comparison with the SSRI in Panic Disorder.

Results

The results are indicated in Figure 15. Figure 15 schematically depicts the CGI-severity score: "fore-going" and "add-on" treatment with pipamperon (8 mg/day; bid) and citalopram (20-40 mg/day; bid) in comparison with the SSRI in Panic Disorder. Treatment with the combo pipamperon-citalopram (n = 3). Treatment with paroxetine is according to the Journal of Clinical Psychiatry (2004) 65: 405-413. Treatment with Sertraline is according to the Journal of Clinical Psychiatry (2004) 65: 405-413.

Conclusion

Notably, although a small test group has been used (n = 3), the distribution around the mean is good. It will further be apparent from Figure 15 that the effect of the combo treatment of pipamperon and citalopram is twice as high as the standard treatments with paroxetine or sertraline.

Example 6: POC process for mayor depressive disorder

Concept: Combo of the high selective 5-HT_{2A/D4} antagonist pipamperon with:

- a compound active towards the Amino Acid Transmitter, Peptidergic Transmitter, Adenosine Transmitter, Endocrine and/ or Enzymatic System;
- a fore-going admission during 4 days of pipamperon;
- a dose of pipamperon of 12 mg/day

Objectives: Demonstrating that this combo therapy has:

- the potency of being a treatment standard for depression by having an added value of reducing the total score of the Hamilton Depression Rating Scale – 17 items (HDRS-17) after 8 weeks of therapy with a least 20% more than reached with the conventional known antidepressants, i.e. 60% versus 40%. This stands for an added medium demission of 5 points on the total score of the HDRS-17 and by this will be very highly

significant since the mean difference in all recent clinical trials between placebo and active treatment is 2.5;

- a more sustained therapeutic effect than the conventional mono therapy by preventing significant more relapses during 48 weeks following the acute treatment; and/or
- 5 - a complete neutral safety profile, e.g. there are no more adverse events in the combo therapy than in mono admission of the in the combo used antidepressant compound.

Process: the following different steps were implemented to reach out for these objectives (see also Tables 3 and 4):

- 10 (1) an naturalistic open label study ($n \geq 20$) on a depressive population with a normal variability of medical and psychiatric history, course of depression, earlier and concomitant therapy admitting the golden standard antidepressant citalopram 20-40 mg/day and a dose of 8-12 mg/day of pipamperon in a foregoing, simultaneous or add-on use.
- 15 (2) a 16 weeks placebo controlled randomised four armed study of each 36 patients with a mayor depressive disorder admitting:
 - from day 0: placebo or pipamperon (PIP) 10 mg/day or an active antidepressant compound or the combination of the last two;
 - from day 4: placebo or pipamperon 10 mg/day combined with an active antidepressant compound or an active antidepressant compound without pipamperon.
- 20

By including rigorous control groups (placebo and active comparator; see Tables 3 and 4) this clinical trial is evaluated as a proof of concept of the added value of the combo and the foregoing treatment method since the inclusion/exclusion of:

- 25 - a negative trial, i.e. no significant difference between the placebo and active treatment with the comparator;
- a failed trial, i.e. no significant difference between the active and the studied treatment i.e. the combo.
- (3) an active controlled randomised relapse prevention study following the POC trial during another 36 weeks with three arms of each 36 patients which is formed by:
 - 30 - continuation of the active mono therapy;
 - randomising the patients with a combo therapy in a group with an active mono therapy and with a continuation of the combo treatment.

Claims

1. Use of pipamperon for the preparation of a medicament for treating a disease or disorder with an underlying dysregulation of the emotional functionality, wherein said pipamperon is administered to a patient in a dose ranging between 5 and 15 mg of the active ingredient, and wherein a second compound is administered simultaneously with, separate from or sequential to said pipamperon to augment the therapeutic effect of said second compound.

2. Use according to claim 1, wherein said pipamperon is administered daily at least one day before administering said second compound.

3. Use according to any of claims 1 to 2, wherein said second compound affects the monoaminergic transmitter system.

4. Use according to claim 3, wherein said second compound is selected from the group comprising: 5-HT reuptake enhancer (1), 5-HT₁ autoreceptor agonist (2), 5HT_{1A} receptor agonist (3), 5-HT_{1A} receptor antagonist (4), 5-HT_{1B} receptor antagonist (5), 5-HT_{2B} receptor antagonist (6), 5-HT_{2C} receptor antagonist (7), 5-HT₃ receptor antagonist (8), 5-HT₆ receptor antagonist (9), adrenergic transmitter releaser (12), α ₁ adrenoreceptor antagonist (13), α ₂ adrenoreceptor antagonist (14), β ₃ adrenoreceptor agonist (19), cannaboid receptor antagonist (21), D₁ receptor agonist (27), D₂ receptor antagonist (28), D₃ receptor antagonist (29), DA uptake inhibitor (30), dopamine receptor agonist (31), H₃ receptor antagonist (42), compounds which increase brain concentrations of 5-HT (44), levodopa (48), MAO reuptake inhibitor (50), MAO-A & MAO-B reuptake inhibitor (51), MAO-B inhibitor (52), MAO-B re-uptake inhibitor (53), NARI (60), NaSSA (61), NDRI (62), RIMA (82), SDA (84), SDRI (85), Second messenger beta agonist (86), SNDRI (90), SNRI (91) and SSRI (92).

5. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorder, personality disorders, antisocial behaviour, bereavement,

occupational problem and problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT (serotonin) reuptake enhancer compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT (serotonin) reuptake enhancer compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

6. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT₁ autoreceptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT₁ autoreceptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

7. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-induced persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment

disorder, other cognitive disorders and Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{1A} (serotonin 1A) receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{1A} (serotonin) 1A receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

8. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{1A} receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{1A} receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

9. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{1B} receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{1B} receptor antagonist compound, further characterized

in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

10. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation
5 of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental
disease or disorder selected from the group of diseases and disorders consisting of mood
disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform
disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and
10 gender disorders, sleep disorders, adjustment disorders, impulse control disorders,
substance-related disorder, personality disorders, antisocial behaviour, bereavement,
occupational problem, problems related to abuse or neglect and pain disorders
characterized in that pipamperon or said pharmaceutically acceptable salt thereof is
administered simultaneously with, separate from or prior to the administration of a 5-HT_{2B}
15 receptor antagonist compound to augment the therapeutic effect or to provide a faster onset
of the therapeutic effect of said 5-HT_{2B} receptor antagonist compound, further characterized
in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and
15 mg of the active ingredient.

11. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation
20 of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental
disease or disorder selected from the group of diseases and disorders consisting of mood
disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform
disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and
gender disorders, sleep disorders, adjustment disorders, impulse control disorders,
25 substance-related disorder, personality disorders, antisocial behaviour, bereavement,
occupational problem, problems related to abuse or neglect and pain disorders,
characterized in that pipamperon or said pharmaceutically acceptable salt thereof is
administered simultaneously with, separate from or prior to the administration of a 5-HT_{2C}
30 receptor antagonist compound to augment the therapeutic effect or to provide a faster onset
of the therapeutic effect of said 5-HT_{2C} receptor antagonist compound, further characterized
in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and
15 mg of the active ingredient.

12. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation
35 of a medicament for treating the underlying emotion dysregulation of substance-related
disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof

is administered simultaneously with, separate from or prior to the administration of a 5-HT₃ receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT₃ receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

13. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT₆ receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT₆ receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

14. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an adrenergic transmitter releaser compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said adrenergic transmitter releaser compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

15. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a alpha 1 adrenoreceptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said alpha 1 adrenoreceptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

16. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a alpha 2 adrenoreceptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said alpha 2 adrenoreceptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

17. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem,

problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a beta 3 adrenoreceptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said beta 3 adrenoreceptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

18. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, pain disorders and delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a cannabiod receptor 1 antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said cannabiod receptor 1 antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

19. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of substance related disorders and Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D1 receptor receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D1 receptor receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

20. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting mood

disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, pain disorders and delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D2 receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D2 receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

21. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D3 receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D3 receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

22. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of substance related disorders and Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a DA (dopamine) uptake inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said DA (dopamine) uptake

inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

23. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders and problems related to abuse or neglect, pain disorder and Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a dopamine receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said dopamine receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

24. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a histamine H3receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said histamine H3receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

25. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform

disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorder, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which increases brain concentrations of 5-HT (serotonin) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which increases brain concentrations of 5-HT (serotonin), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

26. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a levodopa/decarboxylase inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said levodopa/decarboxylase inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

27. A pharmaceutical composition comprising (a) pipamperon, and (b) a levodopa/decarboxylase inhibitor compound, preferably is (eti)levodopa/carbidopa, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof in combination with entacapone, which is an inhibitor of catechol-O-methyltransferase (COMT), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson Disease.

28. A pharmaceutical composition according to claim 27, wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said levodopa/decarboxylase inhibitor compound is levodopa/carbidopa, preferably provided in a unitary dose of between 100 mg and 10 mg of the active ingredient.

29. A pharmaceutical composition according to claim 27 or 28, wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said levodopa/decarboxylase inhibitor compound is levodopa/carbidopa or etilevodopa/ carbidopa

in combination with entacapone, of which the latter is preferably provided in a unitary dose of between 500 mg and 100 mg of the active ingredient.

30. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of non-cognitive mental disease or disorder which are substance related disorders and Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase (MAO) reuptake inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said monoamine oxidase (MAO) reuptake inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

31. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorder, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase A (MAO-A) and a monoamine oxidase B (MAO-B) reuptake inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said monoamine oxidase A (MAO-A) and a monoamine oxidase B (MAO-B) reuptake inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

32. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, problems related to abuse or neglect, pain disorder and Parkinson

disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase B (MAO-B) inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said monoamine oxidase B (MAO-B) inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

33. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase B (MAO-B) reuptake inhibitor to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said monoamine oxidase B (MAO-B) reuptake inhibitor, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

34. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, adjustment disorders, attention-deficit disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorder, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective nor-adrenaline re-uptake inhibitor (NARI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective nor-adrenaline re-uptake inhibitor (NARI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

35. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems

related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a noradrenergic/specific serotonergic antidepressant (NaSSA) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said noradrenergic/specific serotonergic antidepressant (NaSSA) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

36. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, adjustment disorders, attention-deficit disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

37. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which is a reversible inhibitor of mono-amine oxydase A (RIMA) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which is a reversible inhibitor of mono-amine oxydase A (RIMA), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

38. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorder and delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a serotonin/dopamine antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said serotonin/dopamine antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

39. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin and dopamine re-uptake inhibitor (SDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin and dopamine re-uptake inhibitor (SDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

40. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a second messenger beta agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said second messenger beta agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

41. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

42. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep

disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

43. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin re-uptake inhibitor (SSRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin re-uptake inhibitor (SSRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

44. Use according to claim 1 or 2, wherein said disease or disorder is Alzheimer disease.

45. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{1A} (serotonin 1A receptor) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{1A} (serotonin 1A receptor) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

46. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT6 antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT6 antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

47. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an acetylcholinesterase inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said acetylcholinesterase inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

48. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an AMPA receptor mediator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said AMPA receptor mediator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

49. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an amyloid aggregation-inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said amyloid aggregation-inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

50. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a calcium channel modulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said calcium channel modulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

10

51. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a choline uptake enhancer compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said choline uptake enhancer compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

15

52. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound that activates ERK (extracellular signal-related kinase) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound that activates ERK (extracellular signal-related kinase), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

25

53. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a GABA agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GABA agonist compound, further characterized in that pipamperon

35

is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

54. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a histamine H3-receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said histamine H3-receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

55. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which increases insulin sensitivity to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which increases insulin sensitivity, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

56. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which mimics the effect of nerve growth factor (NGF) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which mimics the effect of nerve growth factor (NGF), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

57. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a muscarinic receptor partial agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said muscarinic receptor partial agonist compound,

further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

58. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

59. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from nerve growth factor (NGF) gene therapy, to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said nerve growth factor (NGF) gene therapy, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

60. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a nicotinic receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said nicotinic receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

61. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an N-Methyl-D-aspartate (NMDA) antagonist compound to augment the therapeutic effect or to

provide a faster onset of the therapeutic effect of said N-Methyl-D-aspartate (NMDA) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5 62. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a non-steroidal anti-inflammatory drug to augment the therapeutic effect or to provide a faster
10 onset of the therapeutic effect of said a non-steroidal anti-inflammatory drug, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

15 63. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a phosphodiesterase-4 (PDE4) inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said phosphodiesterase-4 (PDE4) inhibitor
20 compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

25 64. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a peptidic compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said peptidic compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active
30 ingredient.

35 65. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which protects dopaminergic and cholinergic neurons to augment the therapeutic effect or to

provide a faster onset of the therapeutic effect of said compound which protects dopaminergic and cholinergic neurons, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5

10

66. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin and dopamine reuptake inhibitor (SDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin and dopamine reuptake inhibitor (SDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

15

20

67. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin and dopamine reuptake inhibitor (SDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin and dopamine reuptake inhibitor (SDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

25

30

68. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

35

69. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a CRF1 antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said CRF1 antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

70. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a GR antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GR antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

71. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said

pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a melatonin receptor (MT) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said melatonin receptor (MT) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

72. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neurotensin receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurotensin receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

73. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neurokinin 2 receptor (NK2) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurokinin 2 receptor (NK2) antagonist compound, further characterized in that pipamperon

is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

74. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neurokinin 3 receptor (NK3) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurokinin 3 receptor (NK3) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

75. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a peptidic compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said peptidic compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

76. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood

disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a substance P receptor (NK1) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said substance P receptor (NK1) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

77. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a tachykinin antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said tachykinin antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

78. Use according to any of claims 1 to 4 or 7, wherein said second compound is gepirone.

79. A method for preparing a compound having a selective D4 and 5-HT2A antagonist, reverse agonist or partial agonist activity comprising the following steps: (a) measuring the selective affinity of a test compound to the D4 receptor and selecting a compound that has a pKi value equal to or greater than 8 towards the D4 receptor in respect to all the other D receptors, and measuring the selective efficacy of the selected compound to the D4 receptor and selecting a compounds which is a selective antagonist, inverse agonist or partial agonist of the D4 receptor; (b) measuring the selective affinity of a test compound to the 5-HT2A

231


receptor and selecting a compound that has a pKi value equal to or greater than 8 towards the 5-HT_{2A} receptor in respect to all the other 5HT receptors, and measuring the selective efficacy of the selected compound to the 5-HT_{2A} receptor and selecting a compounds which is a selective antagonist, inverse agonist or partial agonist of the 5-HT_{2A} receptor; (c) identifying a compound which is selected in (a) and (b), (d) preparing the compound identified in (c).

80. Compound prepared by the method of claim 79.

81. Use according to any of claims 1, 2 or 7, wherein said second compound is chosen from the group consisting of fluvoxamine controlled release, phenserine tartrate, atomoxetine hydrochloride, bupropion (controlled-release formulation), ropinirole HCL (controlled-release formulation), INN 00835, galantamine (extended release formulation), paliperidone, tomoxetine, aprepitant, rivastigmine tartrate, ORG 34517/34850, sunepitron, sumanirole, milnacipran, idazoxan, xaliproden, SR 58611, befloxatone, litoxetine, tianeptine, agomelatine, SPD 503, flesinoxan, bifeprunox, ramelteon, etilevodopa, rasagiline (TVP-1012) and desvenlafaxine.

82. Use according to any of claims 1, 2 or 7, wherein said second compound is chosen from the group consisting of galantamine (extended release formulation), R121919, risperidone, paliperidone and R228060 (YKP-10A).

**CGI-SEVERITY SCORE: Foregoing & Add-On Treatment with Pipamperon
8 mg/day (bid) and Citalopram 20-40 mg/day (bid) in comparison with the
SSRI in Panic Disorder**




○ - Paroxetine in the treatment of panic disorder, Journal of Clinical Psychiatry 65: 405-413, No. 3, Mar 2004

* - COMBO PIPAMPERON-CITALOPRAM (n=3)

○ - Sertraline in the treatment of panic disorder, Journal of Clinical Psychiatry 65: 405-413, No. 3, Mar 2004


**Y-BOCS COMPULSION SCORE: Foregoing & Add-On Treatment
with Pipamperon 8-16 mg/day (bid) and Citalopram 30-80 mg/day
(bid) in comparison with the SSRI fluvoxamine in OCD**




—○— COMBO PIPAMPERON-CITALOPRAM (n=7)

FLUVOXAMINE Controlled Release mean
271 mg/day (n=253) (Hollander E, Journal of
Clinical Psychiatry 64: 640-647, Jun 2003)

**Y-BOCS OBSESSION SCORE: Foregoing & Add-On Treatment
with Pipamperon 8-16 mg/day (bid) and Citalopram 30-80 mg/day
(bid) in comparison with the SSRI fluvoxamine in OCD**




**Y-BOCS TOTALSCORE: Foregoing & Add-On Treatment with
Pipamperon 8-16 mg/day (bid) and Citalopram 30-80 mg/day (bid)
in comparison with the SSRI fluvoxamine in OCD**




—○— COMBO PIPAMPERON-CITALOPRAM (n=7)

FLUVOXAMINE Controlled Release mean
271 mg/day (n=253) (Hollander E, Journal of
Clinical Psychiatry 64: 640-647, Jun 2003)


**REMISSION RATES (HDRS-17 ≤ 7): Foregoing & Add-On
Treatment with Pipamperon 8-12 mg/day (bid) and
Citalopram 20-40 mg/day (bid) in comparison with the
SNRI venlafaxine in MAJOR DEPRESSION**



HDRS-17 CHANGE FROM BASELINE: Foregoing & Add-On Treatment with Pipamperon 8-12 mg/day (bid) and Citalopram 20-40 mg/day (bid) in comparison with the SNRI duloxetine in MAJOR DEPRESSION




Foregoing & Add-On Treatment with Pipamperon 8-12 mg/day (bid) and Citalopram 20-40 mg/day (bid) in MDD in Comparison with the Standard Efficacy of Antidepressants in Clinical Trials*




* A. KHAN et al, Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical

Trials, ARCH. OF GENERAL PSYCHIATRY / VOL 57, APR 2000)

HDRS-17 CHANGE FROM BASELINE: COMBO PIPAMPERON-
CITALOPRAM WITH A FORE-GOING TREATMENT OF 7 DAYS
WITH PIPAMPERON 11mg/day vs SNRI (duloxetine) in MAJOR
DEPRESSION




Foregoing Treatment During 6-8 (mean 7) days with Pipamperon 8-12 (mean 11) mg/day (bid) Followed With the Combination Treatment of Pipamperon and Citalopram 20-40 (mean 30) mg/day (bid) (PICIT FG 6-8) in MDD (HDRS-17 at BL = 28 in Comparison with the Standard Efficacy of Antidepressants in Clinical Trials*




*** A. KHAN et al, Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical Trials, ARCH. OF GENERAL PSYCHIATRY / VOL 57, APR 2000)**


**REMISSION RATES (HDRS-17 ≤ 7): COMBO PIPAMPERON-CITALOPRAM
WITH A FORE-GOING TREATMENT OF 4 DAYS WITH PIPAMPERON 10mg/day
vs SNRI (venlafaxine) in MAJOR DEPRESSION**



**HDRS-17 CHANGE FROM BASELINE: COMBO PIPAMPERON-
CITALOPRAM WITH A FORE-GOING TREATMENT OF 4 DAYS
WITH PIPAMPERON 10mg/day vs SNRI (duloxetine) in MAJOR
DEPRESSION**




Foregoing Treatment During 1-5 (mean 4) days with Pipamperon 8-12 (mean 10) mg/day (bid) Followed With the Combination Treatment of Pipamperon and Citalopram 20-50 (mean 26) mg/day (bid) (PICIT FG 1-5) in MDD (HDRS-17 at BL = 23) in Comparison with the Standard Efficacy of Antidepressants in Clinical Trials*



* A. KHAN et al, Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical Trials, ARCH. OF GENERAL PSYCHIATRY / VOL 57, APR 2000)

REMISSION RATES (HDRS-17 ≤ 7): COMBO PIPAMPERON AS ADD-ON - CITALOPRAM vs SNRI (venlafaxine) vs SSRI's vs PLACEBO in MAJOR DEPRESSION




— COMBO PIPAMPERON AS ADD-ON - CITALOPRAM WITH PIPAMPERON 8-12mg/day (mean 9mg/day) 20-60 (mean 33) days after the treatment with the SSRI (n=5)
SNRI Venlafaxine - Meta-analysis of Thase et al, Br J Psychiatry 2001; 178: 234-241

— SSRI's - Meta-analysis of Thase et al, Br J Psychiatry 2001; 178: 234-241

— PLACEBO - Meta-analysis of Thase et al, Br J Psychiatry 2001; 178: 234-241


HDRS-17 CHANGE FROM BASELINE: COMBO PIPAMPERON AS ADD-ON - CITALOPRAM vs SNRI (duloxetine) in MAJOR DEPRESSION



COMBO PIPAMPERON AS ADD-ON -
CITALOPRAM WITH PIPAMPERON 8-
12mg/day (mean 9mg/day) 20-60
(mean 33) days after the treatment with
the SSRI (n=5)

DULOXETINE 40-120 mg/day (n=152)
(Goldstein et al, J. Clin. Psychiatry (in
press))

Add-On Treatment with Pipamperon 8-12 (mean 9) mg (bid) after Treatment with Citalopram 10-20 (mean 30) mg (bid) during 20-60 (mean 33) days (PICIT ADD-ON) with HDRS-17 Totalscore = 29 at Baseline in MDD in Comparison with the Standard Efficacy of Antidepressants in Clinical Trials*



* A. KHAN et al, Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical

Trials, ARCH. OF GENERAL PSYCHIATRY / VOL 57, APR 2000)

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/BE04/000172

International filing date: 02 December 2004 (02.12.2004)

Document type: Certified copy of priority document

Document details: Country/Office: CA
Number: 2,487,529
Filing date: 15 November 2004 (15.11.2004)

Date of receipt at the International Bureau: 11 July 2005 (11.07.2005)

Remark: Priority document submitted or transmitted to the International Bureau but not in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☒ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.